

ALPHABET OF INSECTS.

A New Edition.

BY

JAMES RENNIE, M.A.



PRICE HALF-A-CROWN.

ABS. 1.82.186

Sold by
GEO. A. TULLIS.
STATISTICAL ASSISTANT
to the Registrar General
of the Registrar General
of the Registrar General



ALPHABET OF INSECTS,

FOR
THE USE OF BEGINNERS.

BY JAMES RENNIE, M.A.

PROFESSOR OF ZOOLOGY, KING'S COLLEGE, LONDON.

A New Edition,
REVISED, WITH NUMEROUS ADDITIONS.



The Queen Butterfly in its various stages.

LONDON:

ORR & SMITH, AMEN CORNER, PATERNOSTER ROW.

MDCCLXXXIV.



LONDON:
BRADBURY AND EVANS, PRINTERS, WHITEFRIARS,
(LATE T. DAVIDSON).

CONTENTS.

	PAGE
PLAN OF THE WORK	xi
M. Audouin's work taken as a basis	xi
Character of Kirby's terms	xii
Visionary theories	xii
Notice of this edition	xv
THE WORD INSECT	1
Divisions of an insect's body	1
Distinguished from other animals	1
The term <i>Entomology</i> of little use	1
Parts of an insect	2
THE SKIN OF INSECTS—	
Resembles the bones of other animals	4
Fancies built upon this resemblance	4
Layers of the skin	5
Colours of the skin	6
Pores of the skin	7
Hair and down on the skin	7
COMPOSITION OF THE SKIN.	
Discovery and chemical properties of chitine	8
Analyses by Odier, Lassaigne, and Rohiquet	8
THE HEAD IN INSECTS	9
Pieces of the head	9
Form of the head	9
Divisions not determined	9
Parts of the head	10
Members of the head	10
The ears, the eyes, and the mouth	10
Junction of the head with the corselet	11
Absurd theory of Lamarck	11
Position and motion of the head	11
Head of the cockchafer	12
THE CORSELET IN INSECTS	12
Divided into three rings	12
Law of proportion	13
The breast plate	13
The breast prop	13
Its uses	14
The flanks, upper and under	15
The wing scale	15

	PAGE	
The horn scale	16	
Divisions of the back of the corselet	17	
The back plate	17	
The fore back plate	18	
The mid back plate	18	
The back scale	18	
The bridle	18	
The hind back plate	18	
Mechanism to aid the muscles and wings	19	
Middle line of partition	20	
Recapitulation	21	
THE ABDOMEN IN INSECTS.		
Its composition	22	
Its rings	22	
Arches of the back and of the belly	22	
Motions of the abdomen	23	
Cord and pulley mechanism	23	
Muscles	23	
MEMBERS OF INSECTS.		
Meaning of the term	25	
Objection to the term <i>Appendage</i>	25	
MEMBERS OF THE HEAD.		
<i>The ears</i>	25	
Their structure	25	
Their insertion and connexion	26	
Their direction	26	
Their length and proportion	27	
Their form	27	
Ears in male and female insects	27	
THE MOUTH IN INSECTS.		
Number of parts and law of proportion	27	
Wild theory of Lamarck	28	
Reverse views of Savigny	28	
Mistakes of English naturalists	28	
Insects' mouths imperfect and "totally useless"	28	
<i>Parts of the mouth in eating insects.</i>	29	
The upper lip	29	
The upper jaws	29	
The under jaws, with their feelers, outer and inner	30	
The under lip, chin, feelers, and tonguelet	31	
<i>Parts of the mouth in lapping insects</i>	33	
Mouth in bees and wasps	33	
<i>Parts of the mouth in sucking insects</i>	35	
Mouth in two-winged flies, gnats, and bugs	36	
MEMBERS OF THE CORSELET.		36
Legs and wings	36	
Law of proportion	37	

	PAGE
THE LEGS AND FEET IN INSECTS.	
Exactly six in number	37
Jointings of each pair	37
<i>Parts of the legs and feet</i>	37
The haunch	38
The thigh	38
The shank	38
The foot	38
The claw	39
<i>Foot-combs and foot-brushes of insects</i>	39
Cleanliness of flies	39
Opinions of Mr. Blackwall and M. Bauer	40
<i>Muscles of the legs and feet</i>	40
Figure of these muscles	41
<i>Standing of insects</i>	42
Position of the legs	42
Pendulum motions of legs	43
<i>Creeping, walking, and running, of insects</i>	43
Slow walking	43
Quick running	33
Oblique walking	44
<i>Suspensory standing and walking of insects</i>	44
Opinions of Power, Hook, and Derham	44
Experiments of Blackwall	44
<i>Swimming of insects</i>	45
Walking on the water	45
Gliding of the whirlwig beetle	46
<i>Hopping and leaping of insects</i>	46
Leaping of the flea	46
Cuckoo-spit frog-hopper	47
THE WINGS IN INSECTS	47
Their number	47
First pair of wings	47
<i>Parts of the wings</i>	48
Wing joint pieces	48
Wing muscles	48
Thirteen pairs in the cockchafer	48
Structure of the wings	49
<i>Disposition of the wing ribs</i>	50
The edge rib and its hook	50
The mid rib	50
The mid rib branch	51
The mid rib branchlets	51
The inner rib	51
The lower rib	51
The small marginal ribs	51
Terms of the circumference of the wing	51

	PAGE
Areas of the wing	51
Upper area	52
Mid area	52
Lower area	52
Meshes of the wing	52
Base meshes	52
Mid meshes	52
Outer meshes	52
<i>Markings of the wings</i>	52
Streak, band, spot, and eyelet of the wings	53
Fringes and tail of the wings	53
<i>Texture of the wings</i>	53
Horny wings of beetles	53
Muscles of the wings	54
Leathery wings of locusts	55
<i>Peculiarities in the wings</i>	55
Winglet	55
The second pair of wings	55
Poisers	56
Wing bridles	56
Clothing of wings	56
<i>Flying of insects</i>	57
Opinions of Chabrier and Straus Dürkheim	57
Hovering of Wasp-flies	57
Dances of gnats	58
MEMBERS OF THE ABDOMEN	59
Tail Tuft	59
Tail tweezers	59
Sugar-louse	60
Ovipositors	61
Sting	61
Cleaning instrument	61
INTERNAL ORGANS OF INSECTS	62
The order in which these may be taken	62
ORGANS OF DIGESTION	62
Length of the organs in various insects	62
Their three layers	63
The haus	63
Fountains of the mouth	63
Silk organs	63
Division of the organs into six parts	63
Organs in two beetles	64
The gullet and its nervous ring	65
The crop or craw	65
Honey bag of bees and butterflies	65
The gizzard	65
Mistake of Swammerdam and Cuvier	66
Gizzard of the sugar louse	66
The stomach	66

	PAGE
Bile vessels	67
Gastric vessels	67
Outlet of the stomach	67
Intestines	67
The chyle gut	68
The small gut and its functions	68
The blind gut	68
The vent gut	68
The vent	68
Offensive discharges—poison	68
Probable course of the chyle	69
The fluid and solid portions	69
Process of nutrition	70
ORGANS OF CIRCULATION.	70
The heart or dorsal vessel	70
Its resemblance and difference compared with the heart of other animals	70
No direct circulation of blood in insects	71
Futile claims to the discovery of a circulation	71
Extent, position, and form of the heart	72
Its structure	72
Its contents	72
Opinions of Audouin, Meckel, and Herold	72
Description of the heart by Straus Dürckheim	73
Circulation of the blood traced	73
Its supposed distribution and imbibition	75
ORGANS OF BREATHING.	
Insects have no lungs or nostrils	75
The spiracles	75
Simple spiracles	76
Their number and position	76
Wings supposed to be spiracles!!!	76
Lips of the spiracles	76
Composite spiracles	77
The air pipes	77
Air pipes in the cockchafer	77
The two windpipes and their branchings	78
Structure of the air-pipes	78
Structure of the air-cells	79
Their ribs	80
Process of breathing	80
ORGANS OF SENSATION.	
Brain and spinal cord in man	80
Main nervous stem in insects	81
The ganglionic system in man	81
A nerve-knot or ganglion described	81
Ganglionic brain in insects	82
Differs from the ganglia and from the brain in man	82
Gullet ring and second ganglion	82
Third ganglion and others	82
Nervous system in the cockchafer	83

	PAGE
Number of ganglia	84
Structure of an insect ganglion	84
Nerves from the ganglia	84
Structure of the nerves	85
Nervous system in the garden beetle	85
Insects have not acute feeling	86
THE SENSES.	
<i>Rationality and instinct.</i>	86
Dr. Virey's theory	86
Illustration from bees and wasps	87
<i>The sense of touch</i>	87
The ears mistaken for organs of touch	87
Opinions of Bonsdorf and Straus Dürkheim	87
The feet, feelers, and hairs as organs of touch	88
<i>The sense of taste</i>	88
Tongue according to Huber	88
Sense delicate, but the organ unknown	88
<i>The sense of smell</i>	88
Opinions of M. Clerck	89
Opinions of Baster and Huber	89
Singular opinion of Dumeril	89
Opinion of Marcel de Serres	89
<i>The sense of hearing</i>	89
The ears often misrepresented	90
Structure of the ears in the cockchafer	90
Experiments of Bonsdorf, Göze, and Christ	91
Experiments of the Author	91
Remarks of Straus Dürkheim	91
<i>The eyes and the sense of vision</i>	92
Structure of simple eyes	92
Structure of compound eyes	93
Singularities in night insects	94
The nerves of the eyes	94
The air-pipes of the eyes	95
Ants not blind	95
ORGANS OF REPRODUCTION.	
Insects differ from snails, and resemble birds	95
Singularities in bees, wasps, and ants	95
Male insects never assist the female	95
Difference of the sexes in size and colour	95
Difference in wandering and stationary habits	96
All female insects have eggs	96
Popular and theoretical errors refuted	96
Important discovery by M. Ehrenberg	97
Insects' eggs proportionably large	97
Eggs traced in the body of the mother	97
Eggs of the cockchafer	97
Eggs of the louse	97
Egg-glue, and its use	97

	PAGE
Errors of the learned and the unlearned	98
Miner, mason, and carpenter insects	98
Curious instrument of female grasshoppers, saw-flies, and gall-flies	99
Parasite or cuckoo flies	99
GROWTH OF INSECTS	100
Abuse of the term <i>Development</i>	100
<i>Metamorphoses</i> or <i>Transmutations</i> more objectionable than "Transformations"	102
EGGS OF INSECTS.	
Form of eggs	100
Eggs of the gnat built into a boat	101
Markings of eggs	102
The shell expansible	102
Insects do not hatch their eggs	102
Proceedings of earwigs and ants	102
Absurd notions about <i>blight</i>	102
Nobody says <i>blight</i> produces oxen or rooks	102
INFANCY OF INSECTS.	103
No single English word applicable	103
Meaning of the term CATERPILLAR	103
Gooseberry and cabbage caterpillars	104
Turnip fly and "worm i' the bud"	104
Meaning of the term GRUB	104
Not so definite as caterpillar	104
Grubs of cockchafers and of weevils	104
Turnip-fly grub and wire worm	105
Meaning of the term MAGGOT	105
Blow-fly and bot maggots	105
Cheesechoppers and crane-fly grubs	105
Mawks and Gentles	105
Blood-worm and its case	106
The Latin term <i>Larva</i> fanciful and objectionable	106
ADOLESCENCE OF INSECTS.	107
Reason for adopting the term	107
How insects differ here from other animals	107
Preparatory nest or cocoon	107
Covering besides the nest	107
Glit covering, and the term <i>chrysalis</i>	108
Butterflies, moths, flies, beetles, and bugs	108
Terms used by Linnæus objectionable	108
ADULT STAGE OF LIFE IN INSECTS.	109
The fanciful Latin term <i>Imago</i> as objectionable as the similar term <i>Type</i>	109
Injury to real knowledge from such fancies	110
Abuse of the term <i>Perfect</i>	110
Absurd modern distinctions between perfect and imper- fect insects	110
Emergence of the adult insect	110
Peculiar appearance after emergence	110

	PAGE
Importance of the air tubes for inflation . . .	111
Adult insects never increase in size . . .	111
Examples from white butterflies and house flies . . .	111
Mosquitoes of Brazil . . .	112
Short life of adult insects . . .	113
 SYSTEMATIC ARRANGEMENT OF INSECTS.	
Use of a system . . .	114
Its logical convenience as a framework for facts . . .	114
Misrepresented by the disciples of Linnæus . . .	114
Systems lead to serious errors . . .	114
The Trinity proved by the Trinarian system . . .	114
Dangerous errors of the natural system in botany . . .	115
Hibernian blunders respecting moths . . .	115
All systems imperfect . . .	115
The terms <i>kingdom</i> , <i>tribe</i> , and <i>family</i> objected to . . .	115
Fashionable systems rejected . . .	116
System of Fabricius unnecessarily difficult . . .	116
System of Linnæus confused . . .	116
System of Baron de Geer partly adopted . . .	116
 CLASSIFICATION OF INSECTS . . .	 117
WINGED INSECTS . . .	117
<i>Four-winged insects</i> . . .	117
I. Scaly-winged insects . . .	117
II. Hairy-winged insects . . .	118
III. Lace-winged insects . . .	118
IV. Yoke-winged insects . . .	118
V. Equal-winged insects . . .	119
<i>Wing-cased insects</i> . . .	119
I. Cross-winged insects . . .	119
II. Leather-winged insects . . .	119
III. Horn-cased insects . . .	120
<i>Two-winged insects</i> . . .	121
I. Pact-winged insects . . .	121
II. Fan-winged insects . . .	121
III. Poisers . . .	122
 WINGLESS INSECTS . . .	 122
I. Leaping insects . . .	122
II. Springing insects . . .	122
III. Blood-suckers . . .	123

PLAN OF THE WORK.

THIS little Work is intended for those who are desirous of beginning the delightful study of INSECTS, or who, having begun, find their progress interrupted by numerous difficulties. Among these, one of the most prominent is, the want of a plain and short outline of some of the leading subjects of inquiry, by way of a basis on which to proceed—a ground plan to work upon in rearing a superstructure of knowledge.

There is no English work answering to these views, and the French work of M. Audouin, though very excellent in its way, is too brief and too technical*. I have therefore sketched the following outline on a plan, which, I trust, will be found useful, taking M. Audouin's book as a sort of foundation, enlarging where I deemed it too brief, adding where I found any thing important omitted, and avoiding, as much as possible, the intermixing of theory and hypothesis with facts.

In the three little volumes, published in the

* *Resumé d'Entomologie*, 18mo, Paris, 1828.

Library of Entertaining Knowledge, entitled INSECT ARCHITECTURE, INSECT TRANSFORMATIONS, and INSECT MISCELLANIES, I have introduced some of the most interesting details and discussions respecting Insects; but the plan of these works precluded my going into a regular elementary enumeration of parts and functions such as I have here given, and which may accordingly be considered introductory to these three volumes.

The general rule of style which I always adopt, is never to use a word derived from the Latin or Greek when I can readily find one of the Saxon origin, not that it is possible to discard those derivative words altogether—the English language would be meagre indeed without such as have long been naturalised and are well understood; but as they are not the basis of our language, we ought, I think, to try to keep up our native words, and not hunt after foreign terms which we do not indispensably want. “Spotless,” for example, I should generally prefer to *immaculate*; “across,” for *transversely*; “horny,” for *corneous*; “forked,” for *furcate*; and so on. Of all vulgarities, pedantic vulgarity is the most offensive.

In giving names to things not previously named, the Greek language, from its facilities in compounding words, is usually resorted to, but this, I think, is in many cases done without necessity, as English, though not so easily compounded, may be often used with equal, if not greater, advantage. The French in these cases gallicize the Greek; but this, except in chemistry and a few other instances, appears to me in such bad

taste, that I hope never to see the practice followed in England. "Whenever a Frenchman," says a shrewd writer, "can get hold of a rag of Greek, he instantly defiles it*." I have therefore admitted few, if any, terms not of English origin in the text, and have consigned the Latin and Greek terms to the notes.

The serious obstruction to knowledge caused by the rage for multiplying newly invented words is strikingly exemplified in the fact, that, of the "fifty-two pieces composing the thorax," (*corselet*,) "Mr. Kirby does not describe much more than twenty, and yet uses about forty different words for them in his nomenclature†." These are the very words of Mr. Mac Leay, the author's most particular friend. Now, with a book having forty newly coined words for twenty things, how, I may justly ask, is it possible that a student can make progress in a science which has enough of difficulty arising from the minuteness of the objects, without thus taxing invention to increase them? "The doubts and difficulties," says Audouin, "which are experienced at every step, in trying to comprehend what pieces Mr. Kirby wishes to describe, sufficiently prove (do they not?) the total deficiency of such a nomenclature, which is based on no philosophical rule." (*Annales des Sciences Nat.* Fevr. 1832, p. 140, *Note*.)

Another leading feature of this little work is the avoiding of the hypothetical theories which are suc-

* Two Hundred and Nine Days, by T. J. Hogg, Esq. i. 226.

† Zoological Journal. v. 177.

cessively following each other in Natural History, and which often vitiate the most valuable observations of ingenious men. It is not a little singular, that, while these theories are in a great measure banished from other sciences, they should now reign as paramount in this department as alchemy and astrology did in the dark ages. I have not here any room for details; but shall merely refer, for proof of my position, to two works which have appeared within twelve months, one published under the patronage of Government, entitled "Zoology of North America," Part II., in the introduction to which it is gravely suggested that the doctrine of the Trinity may be corroborated by a Trinarian system of Animals!!!—the other, professing to relate to insects, entitled "Sphinx Vespi-formis," 8vo, London, 1832, a tissue of the most incomprehensive fancies I ever looked into*.

The standard works upon insects, for those who study philosophically, and look upon system as only a convenient aid subordinate to their inquiries, are those of Aristotle, Redi, Malpighi, Valisnieri, Swammerdam, Réaumur, Lyonnet, Lesser, Ray, Gould, Bonnet, Leewenhoeck, De Geer, the Hubers, Chabrier, Romdohr, Audouin, Herold, Leon Dufour, Straus-Dürckheim, and a few others, chiefly foreign, our English naturalists being almost exclusively engaged in the inferior departments of classing, inventing terms, and theorizing.

The illustrations have been taken from the best

* See other examples, ALPHABET OF BOTANY, 2d edition, page 153; and FIELD NATURALIST'S MAG., Dec. 1833, page 550.

authorities—Audouin, Lyonnet, Straus-Dürckheim, &c. Those of the parts of the corselet have been reduced from the Zoological Journal, No. 18, by the liberal permission of the publisher, G. B. Sowerby, F. L. S.

Lee, Kent, June 10th, 1832.

NOTICE OF THIS EDITION.

THE present edition has been very carefully revised and numerous additions, and many new engravings have been introduced, chiefly from Straus-Dürckheim, Treviranus, Burmeister, and Bonsdorf. The sketch of the *Systematic Classification* of insects has also been remodelled and enlarged, so as to render it more uniform with the classification in the ALPHABET OF ZOOLOGY. It having been alleged, that, in the first edition, I showed great ignorance respecting the circulation of the blood, and the sense of hearing in insects, I have now given, in their places, a few brief references to some of the highest authorities, which prove such allegations to have proceeded either from a wish to find fault, or from utter novices in the science.

Lee, Kent, 21st Dec. 1833.

ALPHABET OF INSECTS.

THE WORD INSECT.

THE first thing to be learned on beginning to study the branch of Natural History of which I shall here treat, is to know how an insect differs from other animals.

The word *Insect*¹ means "cut into," and every insect accordingly appears with divisions, as if cuts had been made in the body; one just behind the head, and another behind the corselet, as may be seen in a bee or a fly; and the whole body besides is composed of rings, and it is on this account only that modern writers use the term insect.

These cuts divide the body of an insect into three parts, the head, the corselet, and the abdomen; and any animal in which these parts are not found distinct is not an insect. Consequently a spider, a woodlouse, a shrimp, or a crab, is not an insect, because none of

(1) In Latin, *Insecta*; in Greek, *Εντομα*, by Aristotle, whence *Entomology*, a learned term, meaning "the Study of Insects," which I dislike, and very rarely have found it necessary to use.

these have more than two of the divisions; while the earthworm, the leech, the snail, the slug, and the oyster, are not insects, because they have none of these divisions distinct.



Parts of an Insect shown in the African Cricket.—*a*, the head carrying the ears, *f, f*, with the eyes at their base; *b, c, d*, the corselet separated into three pieces; the fore corselet, *b*, carrying the fore or first pair of legs, *i, i*; the mid corselet, *c*, carrying the second pair of legs, *h, h*, and the first pair of wings, *g, g*; the hind corselet, *d*, carrying the hind legs, *k, k*; and *l, l*, the second pair of wings; *e*, the abdomen.

We rank among insects, however, caterpillars, grubs, maggots, gentles, mauks, mealworms, bloodworms, and wireworms; but from these being in a state of infancy or youth, they do not in general possess the three divisions distinct, as they all do when arrived at their last state of full grown or adult insects.

With respect to other marks of distinction, it may be proper to mention that insects have no bones, no veins, no branched arteries, consequently no circulating blood; and do not breathe by the mouth, but through air-pipes in the sides. All insects have exactly six legs, though some butterflies have the first pair so short as not to be readily observed, hence objectionably termed *imperfect* or *spurious* legs by theorists; and many caterpillars have, besides their six legs, as many as ten to sixteen members similar to legs, for the purpose of clinging and climbing.

THE SKIN, OR CRUST, OF INSECTS.

It seems indispensable to every plant and every animal to have a skin, the outer portion of which has no feeling, to cover and protect the more sensible parts in the interior of the body. The skin¹ in insects, however, appears still more important, from its having to fulfil in some degree the office of the bones in other animals.

This office of the skin in insects has given rise to a systematic tissue of fanciful and far-fetched analogies. It having been remarked by Aristotle, and repeated by Willis, with regard to crabs and lobsters, that instead of the flesh covering the bones, the bones cover the flesh, Baron de Geer applied the ingenious fancy (for it is nothing more) to insects, and this led M. De Blainville to arrange all animals into—first, those having internal bones, particularly a spine²; and secondly, those having no internal bones³. M. Geoffroy St. Hilaire carried the same analogies to a very ludicrous length, supposing, for example, that other parts of insects besides the bones have travelled to the outside of the body, such as the lungs, which he represents as having in insects become wings⁴. Even his fancies, however, are tame in comparison with the absurd mysticism of Oken, Carus, and other German writers⁵.

The skin in insects is usually hard, like horn, or tough, like leather or parchment; though in some

(1) In French, *Peau solide*, by Straus-Dürckheim.

(2) In Latin, *Vertebrata*.

(3) In Latin, *Invertebrata*.

(4) See Montagu's *Ornith. Dictionary*, second edition, p. xlvii.

(5) See *ALPHABET OF BOTANY*, second edition, page 154.

species, and also in the first stage of a very considerable number, it is thinner and softer than the human scarf skin.

In all cases the skin furnishes a series of levers, or points of attachment and support to the muscles, by which every motion of the insect must be made. Unlike the skin of animals, therefore, that of insects is made up of various pieces, more or less closely jointed or joined like the various bones of animals. Each bone of other animals, moreover, is well known by a distinct name; but the pieces of the skin in insects have only been recently examined, and the few names already given to the pieces are not well established, and still in some confusion, as has been mentioned above.

An excellent French naturalist, M. Audouin, thinks he has reason to conclude that there is always, from the moment of exclusion from the egg, a determinate number of pieces in every insect, probably thirteen, which are either distinct or two or more of them united; that various species differ in having some of the pieces large and others small, or altogether wanting; and that when any piece is found larger, other pieces near it will be proportionably smaller.

The human skin is formed of three layers, the scarf skin¹, the mucous net work², and the inner skin³. In insects these three parts are also found, according to Straus-Dürckheim, but the coloured portion, which is always the middle layer in man and quadrupeds, is not always so in insects. The colouring matter⁴ in insects is found in two states, one which will dissolve

(1) Technically, *Cuticle*; in Latin, *Cuticula*; in Greek, *Emidionus*.

(2) In Latin, *rete mucosum*.

(3) In Latin, *Corium*; objectionably, *Cutis vera*.

(4) In Latin, *Tela colorata*.

in spirits of wine, and one which will not. That which is soluble is a sort of varnish¹ thinly encrusting the exterior of beetles, and imparting to them brilliant metallic tints, as in the diamond weevil of Brazil. It can only be detached from the scarf skin beneath it by rubbing or scraping, and is never procured in flakes. The insoluble colouring matter is usually brown or black, and never green, blue, yellow, or red; it is not, like the other, found on the exterior, but is contained in the texture of the scarf skin and the inner skin, particularly the first. In the lively coloured dragon flies and grasshoppers, both the scarf skin and the inner skin are as transparent as glass, the colours being produced by a thick coating of colouring matter, like liquid paint, on the interior surface. In those parts of the body which are dull, the insoluble colouring matter is diffused through the texture of the skin. The colouring matter is wanting at the joints.

The scarf skin in beetles is thin, hard, brittle, even friable, and without any trace of fibres. It may in most cases, particularly in the more solid parts, be easily separated from the inner skin beneath, and appears then as a black or brown horny pellicle, more dense in texture than the inner skin.

The inner skin is composed of from three to five layers, which can be separated from each other. These layers are all composed of solid, interlacing fibres, most apparent in the innermost layers. Its colour is not so dark as that of the scarf skin.

The colours of insects are exceedingly various; black prevails among beetles and flies; white among caterpillars, grubs, and maggots, which feed under cover; green among caterpillars which feed on leaves;

(1) In Latin, *Politum*.

grey and dusky among moths; while the colours of butterflies are usually bright white, yellow, blue, red, and black, in diversified shades and markings.

The terms usually given to the colours of insects in our modern English books pretending to science, often exhibit the most offensively pedantic taste, besides being frequently unintelligible. Thus we meet with *griseous*, for grey; *ochraceous*, for buff; *luteous*, for yellow; *miniatous*, for scarlet; *niveous*, for snow white; and numerous expressions of the same kind, which would not be tolerated in any other science.

In many insects an incalculable number of pores may be detected in the skin, and probably these exist in all.

A considerable number of insects are clothed with hair or down issuing through the pores. M. Straus Dürkheim did not succeed in tracing these hairs to their roots; but at the joints they may be observed to arise from two bulbs, one cup-shaped, containing the other, which is a round disc with the hair in the centre.

The hair seems useful in keeping bees warm; in preventing the water from soaking into water beetles; and may also possess electrical uses which we cannot trace. It was proved by M. Odier that the hair of insects differs from that of quadrupeds in chemical composition.

COMPOSITION OF THE SKIN.

IF the student of insects understand any thing of chemistry, it may be useful to mention that the skin of insects is very complicated in its composition. M. Odier, on analysing the wing cases of the cockchafer,

found—1. Albumen; 2. Extractive, soluble in water; 3. A brown substance, soluble in potass and insoluble in alcohol; 4. A coloured oil, soluble in alcohol; 5. Carbonate of potass, phosphate of lime, and phosphate of iron; 6. A particular principle, constituting one-fourth of the weight of the wing cases¹.

According to the analysis of M. Lassaigne, the same wing cases of the cockchafer contain—1. An animal matter, similar to the peculiar matter² of cochineal; 2. A particular principle, the same as that found by M. Odier, constituting two-fifths of the weight of the wing cases; 3. A substance soluble in spirits of wine and in ether; 4. Phosphate of lime; 5. Phosphate of magnesia.

M. Robiquet found that the wing cases of the blister beetle consist of the same particular principle as those of the cockchafer, mixed with a green oil.

It is this principle which in reality forms the outer frame or crust of insects. If an insect, such as a cockchafer or a dung beetle, be plunged into a solution of potass and exposed to heat, the crust of the insect is not dissolved and does not change its form. The only change it suffers is the loss of colour; all the muscles and intestines disappearing, the crust, consisting of chitine, alone remaining.

The substance in question differs therefore from horn, hair, and the human scarf skin, which are soluble in potass. It is besides soluble in hot sulphuric acid, does not become yellow by the application of nitric acid, and burns without fusion or losing its form. It does not contain nitrogen. It consequently resembles woody fibre more than any animal substance.

(1) Technically, *Chitine*, by M. Odier; *Entomeiline*, by M. Lassaigne.

(2) Technically, *Coccine*.

THE HEAD.

THE head ¹ of insects, though it be represented by the new school as composed of three or four pieces, does not in general show the least trace of divisions or rings, like the rest of the body: but, because the rest of the body is composed of rings or pieces, it is conjectured by theory that the head is also so composed, though we cannot perceive the divisions.

In the newly-hatched insect, the head is often joined to the body, as in spiders, without any obvious division, which is always apparent in the adult insect; but, even in the young state, it is usually harder, smoother, and in the grubs of beetles and the young of some other insects, is of a different colour from the body.

In animals which have a bony skull the head is always more or less round; this being both the most capacious form, and the most difficult to injure by accidents ²,—circumstances so important in protecting the brain; but in insects which have no brain, the head, though in most cases round, is often of various shapes; as somewhat square and angular in the stag-beetle, flat and wedge-like in wasps and bees, and nearly triangular in some of the bugs. In the weevil and scorpion fly, again, it is long and pointed, while in some beetles and four-winged flies it tapers backwards into a long neck ³.

Minute naturalists are not yet agreed about the divisions of the head, and M. Audouin first says, "it can be demonstrated that it is composed of several segments;" and again he says, "it is composed of

(1) In Latin, *Caput*.

(2) See ALPHABET OF PHYSICS, page 62.

(3) In Latin, *Collum*.

rather solid walls, most frequently presenting no trace of junction, so that it appears at first view quite simple, though an experienced eye soon discovers it to be the result of several segments—the number not yet determined—united together." M. Straus-Dürckheim, in the same vein, says that though the head seems to be composed of one piece, yet on studying it in detail, and comparing it with the head of a centipede, it is easily determined to consist of several segments united into one mass.

That is, as it appears to me, a theorist, accustomed to find that the bones of a skull can be separated, readily imagines junctions of pieces in the head of an insect, though he acknowledges them to be imperceptible.

The head of insects, like that of other animals, consists of the crown and forehead (both of which M. Straus-Dürckheim and Meckel objectionably call the skull¹), and the face; but the extent of the cheeks is seldom very distinct. It is not so with the mask², as we may call it, a part which on the one side is jointed with the front or face, and on the other with the upper lip, which it covers; and when the upper lip is wanting (as it is in some species), the mask performs its office.

In all insects, two horn-like members project from the forehead—very long, with many joints, in butterflies; very short in the house fly; and prettily feathered in male gnats, and in many male moths. These I shall venture to call ears³, as it is most probable, as will be afterwards shown, though not distinctly proved, that they are the organs of hearing. In front, and at the

(1) In Latin, *Cranium*.

(2) The mask is termed by the French, *Chaperon*; in Greco-Latin, *Epistoma*.

(3) In Latin, these organs are termed *Antennæ*.

root of the ears, are two eyes; but many insects have besides several other eyes on the forehead, as may be seen in bees, who have three of these frontal eyes, as will be figured in a subsequent page. Insects have no organ similar to the nose in other animals, as they breathe entirely by the air-pipes in the sides. The mouth is, for the same reason, not employed either for breathing or for uttering sounds, but simply for feeding. It is very different in form and structure from the mouth of other animals.

The head is joined to the first ring of the corselet, either by a neck, by a simple membrane, or by means of a socket hollowed into the form of a funnel.

M. Straus-Dürckheim describes a small bridle under the skin, consisting of two pieces, the fore piece much longer than the hind piece, of which beetles have always two pairs. Their office is to unite the head to the corselet; and following the fanciful theory of Lamarck, he farther supposes them to be the *remains*¹ of two rings, whose breast and sides have in the course of ages disappeared, giving as a reason that the breasts and sides have *not yet* disappeared in the earwig and the cockroach².

The natural position of the head is various. In grasshoppers and dragon flies it is vertical; in most beetles it is somewhat slanting, and almost horizontal. It is very moveable in the common fly and the dragon fly, so that it can be turned almost round, as on a pivot. In many other insects it is immoveable, as in the grasshoppers. In others, the head can be folded down, or drawn back, so as to remain concealed, or nearly so.

The head of the cockchafer is described by M.

(1) In French, *Restes*.

(2) See Montagu's Ornith. Dict., second edition, page xli.

Straus-Dürckheim to be moved by two orders of muscles. The elevator, for raising the head, has two parts, and is rather strong; the depressor, for lowering the head, is weak and short; the rotator, for turning the head, has two parts, which raise the head when both act together; and the flexors, which bend the head on one side. The neck bridle is moved by a retractor muscle, by an oblique elevator, and a straight elevator.

THE CORSELET.

THE middle portion of the body in insects, always obviously distinct from the head before, and from the abdomen behind, I shall term the corselet¹. It is similar in situation and somewhat in office to the chest in larger animals. Unlike the head, this part is composed of several distinct rings and pieces, which have but recently received names and been studied. Most of the names invented by our English authors are confused, inappropriate, and bad. In following Audouin and Straus-Dürckheim, I shall endeavour to be as plain and simple as I can.

The most common division of the corselet is into three rings or segments, but though these are sometimes very distinct, they are also sometimes intimately united, as we have seen is the case with the head. These three rings, beginning from the neck and proceeding backwards, may be termed the fore corselet², the mid corselet³, and the hind corselet⁴, which three being united, form the corselet. It will be convenient to consider that the whole corselet, as well as its three rings, has a breast, two sides, and a back.

(1) In Latin, *Thorax*.

(2) In Latin, *Prothorax*.

(3) In Latin, *Mesothorax*.

(4) In Latin, *Metathorax*. M. Straus-Dürckheim restricts the term "corselet" to our fore corselet, and terms the midcorselet, *Prothorax*.

The student must be warned that among the numerous species of insects, these several parts are found to be very variable in size, and no less so in form, as is minutely described by M. Straus-Dürckheim; agreeably to the interesting law, that as one part is found enlarged another is found proportionally diminished, it is not unusual for a particular piece of the corselet to be exceedingly minute, or wholly wanting, if a contiguous part be comparatively very large.

Taking the whole corselet, and beginning at the breast, we find a piece which may be appropriately termed the breast-plate¹, and is found in all insects, forming sometimes a large, sometimes a small piece; in some species very distinct, as in several water-beetles, but in others so intimately united to the contiguous pieces that the joining is imperceptible. The breast-plate forms the middle of the breast in all the three rings of the corselet, and for the sake of more minute distinction, it may be considered as divided into the fore breast-plate², the mid breast-plate³, and the hind breast-plate⁴.

On the inside of the breast-plate, in the interior of the breast, is another singular piece, usually jointed with the hinder and inner end of the breast-plate. It varies much in form, but frequently resembles a Y, and is hence called the prop, or Y piece, by the French naturalists⁵. This is always present in each of the three rings, and, like the breast-plate, may be considered as the fore breast-prop⁶, the mid breast-prop⁷, and the hind breast-prop⁸. It extends in some cases

(1) In Latin, *Sternum*, which is inaccurate.

(2) In Latin, *Prosternum*, or *Sternum anterius*.

(3) In Latin, *Mesosternum*, or *Peristethium*, which is bad.

(4) In Latin, *Metasternum*, or *Mesostethium*.


(5) In Latin, *Furca*, or *Entothorax*.

(6) In Latin, *Antefurca*, or *Stigma primum*.

(7) In Latin, *Profurca*.

(8) In Latin, *Postfurca*.

within the head, and in others, as in the tree-hoppers, within the abdomen, and was then termed by Réaumur the scaly triangle.



The breast-plate in the great water beetle (*Hydrophilus piceus*); A, outer breast-plate; *a, b*, outer breast-prop; *c, c*, the under flanks; *d, d*, hinder wings. B, inner breast-plate; *e, f, g, g*, inner breast-prop.

The chief use of the breast-prop, seems to be to afford an attachment and lever to the muscles which move the fore legs; very evident in the singular breast-prop of the mole cricket: and, according to Audouin and Marcel de Serres, to protect the nerves, by separating them from the gullet, stomach, and heart. The hind breast-prop has often attracted attention from its singular trident-like form, as may be seen in the preceding figure at *f, g, g*.



The fore corselet, under side; *a, b*, the breast-plate; *c, c*, the under flanks.

The side pieces, (Dr. Burmeister calls them shoulder blades,¹) which may be called the flanks, consist of one on each side, in each of the three rings of the corselet, joined to the breast-plate,—in all three pairs, which we may call the under flanks²; and three other pairs, usually united to the former above and behind, sometimes even resting on the breast-plate itself, may be called the upper flanks³. These have always a constant relation with the haunches⁴ of the ring to which they belong, and sometimes share in forming the circumference of their socket, being jointed with them by means of another small piece, which M. Audouin calls *Trochantine*⁵. Dr. Burmeister only found it in water beetles.



The mid breast-plate; α , the breast-plate; b, b , the mid under flanks.

Besides these six pairs of flanks, there is generally, according to M. Audouin, though not always, a very small piece, in relation with the wings and the under flank, always resting on the latter, sometimes going along its fore border, but sometimes, becoming free, it passes either before or above the wing. It may be termed the wing scale⁶. Dr. Burmeister says he does not know the part in the hind breast-plate.

Along the fore border of the under flank, sometimes

(1) In Latin, *Scapula*, which is bad. (2) In Latin, *Episterna*.

(3) In Latin, *Epimera*. (4) In Latin, *Cocœ*.

(5) In Latin, *Rotula*, by Straus-Dürckheim.

(6) In Latin, *Parapter* or *Patagium*, by Burmeister.

of the breast-plate itself, and even of the upper part of the body, the opening of an air-pipe or spiracle may often be observed surrounded by a small piece, often of a horny texture, which may be called the horn scale¹. It is not always present, for the spiracle itself is often closed; but when it is found, it is important, as a good guide in the comparison of the other parts. The scale is moved by a small muscle.

It is proper here to remind the student, that, when he examines an insect, he ought to look for all these pieces, just as, in examining a rat, he ought to look for the gall bladder; but he must not be disappointed if he do not find all the pieces, as some may be wanting, as the gall bladder is in the rat; and the accounting for such variations opens a fine field of philosophical research.



The hind breast-plate; *a*, its upper flank²; *b*, its under flank³; *c*, the plate itself; *d*, the breast-prop.

Having thus gone over the various pieces which compose the breast and the sides of the corselet, we shall next turn to the upper part of it, which may be termed the back-plate⁴, extending from the neck to the abdomen. In reference also to the division into

(1) In Latin, *Peritrema*, or *Stigma primum*.

(2) In Latin, *Ischium secundum*, by Straus-Dürckheim; or *Pleura*, by Burmeister.

(3) In Latin, *Ischium primum* or *Parapleura*.

(4) In Latin, *Tergum thoracis* or *Notum*, by Burmeister.

three rings, and beginning at the head, we may call the first, the fore back plate¹; the second, the mid back plate²; and the third, the hind back plate³.



The back of the whole corselet in a wasp (*Polistes Bil'ardi*). *a*, the fore back plate; *b*, *c*, the mid back plates; *c*, the mid back scale; *e*, *g*, the hind back plate; *e*, the first piece; *f*, the second piece; *g*, the hind back scale; *d*, *d*, the sides of the mid back plate; *h*, the fourth piece of the hind back plate; *i*, *i*, the under flanks; *k*, *k*, the wing scales of the second pair; *l*, *l*, the wing scales of the first pair; *m*, *m*, haunches of the hind legs; *n*, part of the abdomen; *o*, *o*, *p*, *p*, *q*, *q*, the sockets of the two pair of wings; *r*, *r*, the spiracles for breathing; *s*, the cord for the pulley.

Each of these, in many instances, is composed of four pieces. The fore back plate is thus found to consist of four pieces in locusts and crickets, being in these insects large, and the breast small, according to the law of proportion. In beetles, and most winged insects, however, only two pieces can be readily dis-

(1) In Latin, *Tergum prothoracis* or *Pronotum*.

(2) In Latin, *Tergum mesothoracis* or *Mesonotum*.

(3) In Latin, *Tergum metathoracis* or *Metanotum*.

tinguished, on account of the intimate junction. The fore part¹ is generally concealed within the body.



The fore back plate, front and side views; *a*, the first piece, or collar; *b*, the second piece².

The mid back plate is, in like manner, composed of four pieces. The second of these³ is important, from its always jointing, by means of side pieces, (very distinct in the ruby tail fly,) with the wings. The third is what may be called the back scale⁴, and is usually somewhat triangular in form. In some of the plant bugs, this part extends over the wings, wing cases, and the border of the abdomen. Another piece of the mid back plate is usually quite concealed in the body, or sometimes so intimately joined to the former, as scarcely to be distinguishable. Sometimes it is distinct and free. It has been termed the bridle⁵.

The hind back plate is also composed of three pieces, the first being often concealed within the body, but

(1) In Latin, *Præscutum* or *Prophragma*, by Burmeister.

(2) In Latin, *Pubis*, by Straus-Dürckheim, or *Omium*, by Burmeister.

(3) In Latin, *Scutum*.

(4) In Latin, *Scutellum*.

(5) In Latin, *Frænum*.

obvious in bees and wasps. The second is occasionally composed of two pieces, as in some water beetles, but



The mid back plate, front view. *a*, the fore part; *b*, the second piece; *c*, the third piece; *d, d*, the supposed side pieces of the second piece; *e*, the internal third piece; *f, f*, the wing scales of the first pair; *g, g*, joint bones of the first pair of wings above the sockets.

these are intimately united in the stag beetle. The third piece is divided by a channel lengthwise, which has led to the mistake of considering it as two separate pieces. The fourth piece is very conspicuous in the common fly, and also in the cockchafer; but, in obedience to the law of proportion, it is exceedingly small in crickets, grasshoppers, and locusts.

In order to render this brief description of the corselet complete, it may be necessary to mention that, on the inner surface of the plates, there are certain remarkable inequalities of surface, and even some distinct parts observable, besides those already described.

Amongst these are occasional horny ridges¹ always formed by the junction of two contiguous pieces; and when they can be detected, they are excellent guides for discovering the boundaries of two pieces



The hind back plate, front view. *a*, the first piece; *b*, the third piece, the second not being obvious; *c*, the fourth piece; *d, d*, the under flanks; *e, e*, the wing scales of the second pair; *f, f*, the breathing spiracles; *g*, the cord for pulley.

which cannot be distinguished on the outside. They serve for the attachment of muscles. When these ridges are on the outside², they often assist in the mechanism of a joint, such as those of the wings.

Other small pieces are found in the inside of the corselet, sometimes in the form of flat plates raised upon a foot stalk, and wide at top, like certain mushrooms, such as those which Réaumur found in the abdomen of the tree-hopper, and termed cartilaginous plates. Like the former they either serve for the attachment of muscles³, or assist in the mechanism of the wing joints⁴.

It is farther to be remarked, that if the corselet, as

(1) In Latin *Apodema insertionis*.

(2) In Latin *Apodema articulationis*.

(3) In Latin *Epidema insertionis*.

(4) In Latin *Epidema articulationis*.

well as the head and abdomen, be cut exactly in half, lengthwise, each of the pieces and organs on each side will correspond as exactly as our two hands, or our two nostrils; a very remarkable fact.

Recapitulation.

The corselet, then, or chest of an insect, according to this statement, consists of

FORE CORSELET	{	Fore breast plate, without, Fore breast prop, within, Two under flanks, Two upper flanks Fore back plate, four pieces.
MID CORSELET	{	Mid breast plate, without, Mid breast prop, within, Two under flanks, Two upper flanks, Two wing scales, Mid back plate, four pieces.
HIND CORSELET	{	Hind breast plate, without, Hind breast prop, within, Two under flanks, Two upper flanks, Two wing scales, Hind back plate, four pieces.

In all thirty-six pieces; or, if considered as divided lengthwise in the middle, there will be fifty-two pieces. The pieces when moveable have a number of muscles of which four pairs belong to the fore corselet; five pairs to the mid corselet; and what M. Straus Dürckheim conjectures to be a respiratory muscle to the hind corselet. When the pieces are not moveable the muscles serve for bracing them firmly together.

THE ABDOMEN IN INSECTS.

WE are obliged to use this very objectionable term for the third and last division of the body, which is more or less closely united to the corselet. The abdomen may be described to be that part of the body

which succeeds the corselet, consisting in most cases of a certain number of rings, without legs, wings, or other jointed members for locomotion, and uniformly enclosing a portion, sometimes a very small one, of the intestines.

It is formed by a series of very short hollow cylinders or rings, united with each other by a joint, by a membrane, and sometimes by an intimate junction, the exact line of which is not obvious. Sometimes the rings slide into one another like the tubes of a telescope.

Each of these cylinders is called a ring or segment¹ and is sometimes composed of a single piece sometimes of two half cylinders, whose two borders usually come into contact. In other cases they do not touch at this point, but remain free, and one more or less overlaps the other, as in bees.

Each ring is virtually composed of two principal portions, which, when they can be distinguished, (this is not always possible) take the name of arches². The upper is called the arch of the back³; the under, the arch of the belly⁴.

In the flea, the bed bug, and other insects without wings, as well as in grubs and caterpillars, where the joining of the corselet with the abdomen is not so obvious, the latter may always be known by the legs never being jointed with it.

When the back of the abdomen is covered, as in beetles, by the wing-cases, it is softer and more flexible than the belly; in other cases, the reverse.

In beetles, and some other insects, the abdomen is joined to the corselet without any joint to permit mo-

(1) In Latin, *Segmentum*.

(2) In Latin *Arcus*.

(3) In Latin *Arcus tergi*.

(4) In Latin *Arcus ventris*.

tion ; while in bees, wasps, and most two-winged flies, there is a very obvious joint, consisting of a hinge scooped out in the first ring for the purpose of receiving a projecting part of the fourth piece of the hind back plate of the corselet, arising from between the thigh sockets of the third pair of legs.



The termination of the hind back plate ; *a*, joint of the abdomen ; *b*, *b*, sockets of the thighs of the hind legs.

There is besides, in the corselet, a distinct opening or hole, (of a triangular form in bees,) to give passage to the broad tendon¹ of a muscle from the abdomen, and to serve as a pulley², over which it may play. In proportion as this muscle contracts, it accordingly pulls the abdomen upwards, which falls downwards again when it is relaxed.

Wasps, bees, earwigs, and many other species, have the power of moving the abdomen in various directions, as a whole, and of bending and curving it as a dog does his tail, all of which motions are performed by means of muscles attached to the inner surface of the rings.

The muscles or cords by which every motion is performed are very numerous, as may be seen from those of the caterpillar of the goat moth, traced with incredible skill, patience, and accuracy by M. Lyonnet, to whom I am indebted for the following figure.

(1) In Latin, *Funiculus*.

(2) In Latin, *Trochlea*.



Muscles of the caterpillar of the goat moth (*Cossus ligniperda*): with the two main air-pipes running along each side, and the heart with its six pairs of pyramidal wings in the middle.

In the cockchafer, M. Straus Dürckheim describes the second ring as moved by an upper and an under pertractor, the first having two heads; and the third ring as moved by an under pertractor also consisting of two rather distant heads. The other rings have very large upper pertractors, and under pertractors, united at their edges so that their separations are indistinct. The under pertractor of the last ring is very large. The last arch has also an elevator and a depressor.

A row of round openings may be traced along each side of the abdomen, which are the spiracles or pores for breathing. Each spiracle is shut and opened by a very small sphincter muscle.

MEMBERS OF INSECTS.

THE word member has been chosen here from its involving no theoretical fancy, as the term *Appendage*, used in modern books, always seems to do. I apply the term member to any part of an insect, either jointed upon the body, or not appearing to make a portion of its surface, such as the legs, wings, and ears, as well as the eyes; and, by slightly extending the term, it may also include the tail-fork of the earwig, and the sting of the bee.

In describing the members, I shall follow, in the same order as before, the head, corselet, and abdomen.

MEMBERS OF THE HEAD.

IT will be most convenient to consider the several members of the head, in the order in which they appear most conspicuous to the eye, beginning with the ears, and proceeding to the feelers, the lips, and the jaws.

The Ears of Insects.

The members which I here venture to call the ears¹, though not quite proved, as we shall afterwards see, to be the organ of hearing, are uniformly two in number, standing out from the head, upon which they are jointed and moveable in a socket² by means of a ball or pivot³.

Supposing, with Audouin and Straus-Dürckheim, the head to be composed of a ring made up of determinate pieces, having a breast, back, and two upper

(1) In Latin, *Antennæ*, which means placed *before*, and was applied by the Romans to the sail-yard of a ship.

(2) In Latin, *Torulus*.

(3) In Latin, *Bulbus*.

and two under flanks, similar to the corselet, the ears would be found to arise like the wings, from the junction of the upper flanks with the back or rather crown plates. The ears are therefore fancifully alleged to be *repetitions* of the wings by the German school.

The ears are composed of minute cylinders or rings successively added to each other, to the number of thirty in some butterflies; and thus forming a tube, which encloses nerves for sensation and muscles for moving, as well as air-pipes and cells, as we shall see anon.

As to their insertion or connexion, they are always, according to M. Audouin, placed near the eyes, before, behind, above, below, between, or even apparently within these. Sometimes their bases are near together, or united, and sometimes considerably distant.

As to their direction, they are stiff or flexible, straight or nodding, parallel or diverging, spiral or not spiral; and they are in some species carried always forward, in others backward, or towards the sides, and sometimes folded up or drawn into a sheath.



Various forms of the ears of insects.

As to length, they are very long compared with the length of the body, in some moths and beetles, and very short in the house-fly; but their length does not depend on the number of the joints, for they may be long when composed of only three or four pieces, and short when composed of ten or more pieces.

As to form, they are either hair-like (*a*), cylindrical, conical, bristle-shaped, awl-shaped, spindle-shaped, sharp-pointed (*b*), forked, spike-like (*c*), branched, feathered (*l*), thickened (*e*), tiled, beaded (like a necklace) (*f*), pectinated (like a comb) (*k*), serrated (like a saw) (*d*), prism-shaped, whirled (*h*), downy, hairy, bristly. Their tips again are either pointed, knobbed (*g*), clubbed, hooked, triangular, leaved (*m*) (as in the dung beetle); forked, blunt, awned; abrupt, perforated, (*i*); or fan-like.

In the males of moths, gnats, and some other insects, the ears are in all or most cases more ornamented with feathers, hairs, or sculpture than in the female, in which the ears are usually plain.

The ears are moved by a pair of strong flexor muscles behind, a pair of pertractor muscles more slender before, and a pair of elevator muscles as slender as the pertractors. Besides these, each joint of the ears is furnished with a pair of abductors and a pair of adductors, all figured in a subsequent page.

The Mouth in Insects.

Following the generalising views of Audouin and Straus-Dürckheim, in considering the head composed of similar parts to one of the three divisions of the corselet, the mouth will occupy the place of the breast-plate and the two other flanks; and taking each of those three as composed of two pieces, there would be in the mouth, by supposition, six pieces; and this, in

fact, is the number of pieces found in a great number of species, though not in all; for, by the law of proportion, when a part is much enlarged, a contiguous part is either very small, or altogether wanting.

Notwithstanding the mouths of insects are thus composed of a determinate number of pieces, their structure both appears to be, and is, very different; for example, in a bug, a butterfly, a bee, and a beetle, owing to the difference of form, as well as the difference of junction in the several pieces.

M. Lamarck, who seems first to have had a glimpse of the general uniformity in the number of pieces in the mouth of all insects, at once leapt to the singular and untenable conclusion, that bugs, through process of time, got rid of the joinings that made their mouths into a tube, and successively improved themselves into beetles with good moveable jaws. M. Savigny, reversing this process, thinks the jaws of the beetles *degenerated* into the suckers of the butterfly and the bee.

Many of our English naturalists, from being far behind in logic and generalising, and therefore incompetent to take advantage of continental researches, so admirable when they are stript of theory, forthwith conclude that all insects, without free, moveable jaws, or having any of the six pieces wanting, have *imperfect* mouths. One English naturalist in particular, by a gross misconception of Savigny's meaning, represents the mouths of sucking insects as "totally useless;" and thence concludes, "they can do injury to agriculture;" a conclusion as false as the former is impious.

I have stated this in order to prevent misconception, which, from the imperfection of terms, is but too apt to mislead a genuinc field observer, and is certain to mystify and bewilder a compiler or a cabinet naturalist.

Parts of the Mouth in eating¹ Insects.

Comparing the jaws of an insect with the bill of a bird, we find that, while the motion of the bird's mandibles is upwards and downwards, the motions of the insect's is forward from the sides. According to Savigny, Audouin, Straus-Dürckheim, and Burmeister, the mouth² of an insect consists of the upper lip, a pair of upper jaws, a pair of under jaws, and an under lip.

1. The upper lip³ is a flat, usually horny, plate, joined horizontally to the lower part of the face, and closing the mouth.

It is sometimes lobed and has often a large projection above. It is wanting in the stag beetle, its place being supplied by a projection from the forehead. In the sugar-louse⁴ it is triangular and divided into four moveable portions as figured in p. 33 (*a*). There are never any pieces attached to the upper as there are to the under lip. It is moved by muscles, of which there is in the cockchafer a single pair of elevators.

2. The upper jaws⁵, the only incisive instrument in the mouth, one on the right and another on the left, resemble, in eating insects, a large, horny, solid tooth, more or less curved, often indented, and jointed into the sides of the head immediately below the upper lip, being moveable, and without any pieces attached to them, as is the case with the under jaws.

In the sugar louse, there are five teeth towards the tip of each jaw. The upper jaws are used by insects of prey to seize on their quarry, and in such they are long, while in those which feed on vegetables, or carrion, they

(1) In Latin, *Manducantia*.(2) In Latin, *Os*.(3) In Latin, *Labium* by Illiger, *Clypeus* by Fabricius.(4) In Latin, *Lepisma saccharinum*.(5) In Latin, *Mandibulæ*.

are short. The stag beetle which lives on the juice of trees is a remarkable exception; but its long powerful upper jaws are chiefly defensive. The rose chafer which feeds on soft pollen has the tips of the jaws soft. In most beetles, there is a tuft of hair near the base of the jaw, supposed by Straus Dürckheim to be the organ of taste. Each upper jaw has two muscles, an adductor very strong for shutting the jaws, and an abductor more feeble for opening them.

3. The under jaws¹ are also two in number, and are jointed into the right and left of the inner cavity of the mouth, immediately below the upper jaw. They resemble the upper jaws in moving from the sides forward; but are seldom so strong, being rather membranous than horny, particularly at the tips. They also differ in being jointed, while the upper jaws are solid.

In the sugar louse they are bulged behind, and taper off at the tip, where there is a moveable blade like that of a pair of callipers.

M. Straus-Dürckheim describes each under jaw as composed of three pieces, namely, the body, the feeler, and the shield. In the cockchafer, and other beetles, the body consists of four pieces, the hinge², the stem³, the scales⁴, and the lobe⁵.

There are three muscles for moving the wings; the abductor, a small muscle, the pertractor much stronger, and the adductor. The lobe has two muscles, a flexor and an adductor, both very long.

Usually, at the place where the horny texture ends, and the membranous begins, there is, on each of the

(1) In Latin, *Maxilla*. (2) In Latin, *Cardo* or *Ramus transversalis*.

(3) In Latin, *Stipes* or *Pars dorsalis*.

(4) In Latin, *Squama palpifera*.

(5) In Latin, *Lobus* or *Pars intermaxillaris*.

under jaws, a remarkable little member, in form of a thread, composed of from four to six joints, commonly tipped with a part less horny than the rest, jointed upon a stem or footstalk that supports it, and furnished with hairs. These jointed members are called feelers¹, though the term is objectionable, because their use is not well ascertained.

Sometimes the inner front of the lower jaw forms a sort of acute lobe, or even a large scale, furnished at the tip with a hook, and resembling the upper jaw. In that case, the outer division of the upper jaw takes sometimes the form of a case or arched shield²; sometimes it constitutes a second feeler which is short, and inserted within it. These are called inner feelers³, to distinguish them from the outer, and longer feelers⁴, described above. Hence, some beetles appear to have six feelers. The shield has two muscles, an abductor and an adductor; the feelers have each the same.

The only organs in other animals similar to these feelers are the whiskers in the cat, the mouse, the seal, the night-jar, and other birds which feed on insects; and the beardlets in the cod, the barbel, and other fish; and hence, I think whiskers would be a less objectionable name for them than feelers.

4. The under lip⁵, which closes the mouth below, differs much from the upper lip, being not unlike a second pair of upper jaws united, on their inner side, and covered in the greater part by a horny projection called the chin⁶, containing in its cavity muscles for moving the tongue.

Each of the halves of the under lip carries a feeler⁷ smaller than those of the under jaws, and composed of

(1) In Latin, *Palpi* or *Antennulæ*, by Latreille.

(2) In Latin, *Gates*.

(3) In Latin, *Palpi interni*.

(4) In Latin, *Palpi externi*.

(5) In Latin, *Labium*.

(6) In Latin, *Mentum*.

(7) In Latin, *Palpus labialis*.

four or more joints. A projection in front of the chin and jointed upon it is called the tonguelet¹. In many insects there is found a small piece, which may be termed the tongue scale², on each side, resting upon the tonguelet, with a tip like a small ear.

A single pair of elevator muscles moves the entire lip; while the lip feelers are moved by three pairs, elevators, depressors, and abductors.



Parts of the mouth in a beetle. *a*, the upper lip; *b*, the upper jaw; *c*, the under jaws, with two pairs of jointed feelers; *d*, the under lip, with the short tongue in the middle, the chin below this, and a pair of jointed feelers.

(1) In Latin, *Lingula*. (2) In Latin, *Paraglossa*.

Fig. 1.



Fig. 2.



Fig. 3.

Parts of the mouth in the sugar louse magnified. Figure 1, *a*, the upper lip; *c*, base of the upper lip; *f, f*, the ears, with the tips removed; *h, h*, the lip feelers; *M, M*, the upper jaws; *m, m*, indentations thereof; *o, o*, the eyes; *p, p*, the jaw feelers. Figure 2, *a, a*, basal joint of the lip feelers; *b, b*, second joint; *c, c*, third joint; *d, d*, tip joint; *n, n*, base lobes of the under lip; *s, s*, tip lobes. Figure 3, under jaw magnified; *l, k*, the body; *P*, the feeler; *g*, the shield.

Caterpillars and grubs have their mouths formed very similarly to those of beetles.


Parts of the Mouth in Lapping¹ Insects.

In insects, such as bees and wasps, which may be said to lap, the organs of the mouth are considerably different in form and size. If we hold a bee between the fingers, we can perceive a brown shining instrument², curved like a surgeon's needle, folded closely

(1) In Latin, *Lambentia*.

(2) In Latin, *Rostrum*, or *Proboscis*.

down from the mouth towards the throat. This the bee can project forward like the beak of a bird, or can curve at pleasure. By gentle pressure, the seven pieces of which the whole is composed may be brought into view, as in the figures from Burmeister.



Beak of the honey bee magnified. A, upper view; B, under view; *a, a*, the head; *b*, the three simple eyes; *c, c*, the faceted eyes; *d, d*, the ears; *e*, the mask; *f*, the upper lip; *g, g, g, g*, the upper jaws, with their teeth; *h, h, h, h*, the under jaws which, when united, form a sheath for *i, i*, the tongue; *k, k*, jaw feelers, very small; *l, l, l, l*, lip feelers; *m*, the chin; *n*, the tongue prop¹; *o, o*, the tongue scale.

What I have here, after Cuvier and Burmeister, termed the tongue, is by M. Audouin taken for the under lip. He says, the under jaws are rather long,

(1) In Latin, *Fulcrum*, or *Os hyoideum*, which is bad.

encasing the sides of the lip, and, these, being united, form a sort of sucker, moveable at the base.

Parts of the Mouth in Sucking Insects¹.

In butterflies and moths, the upper jaws and the upper lip are very minute; the under jaws are immovable at the base, and form two lines which are united and form a long tube rolled up spirally. It would be no less incorrect to call this the tongue, as it has been called by Fabricius, Latreille, and Cuvier, than it would be to represent it as an *imperfect* mouth, as has been done by our English naturalists.



Parts of the mouth in the green forester², a day flying moth.—*a*, the upper lip; *b, b*, the upper jaws; *c, c*, side view of the under jaws in form of a sucker carrying the jointed feelers at their base; *d*, the under lip with the tonguelet in the middle, and the jointed feelers on each side.

In two-winged flies and gnats, the upper lip forms a case, the two pair of jaws are in form of bristles, and

(1) In Latin, *Suctoria*. (2) In Latin, *Zygana Stylicæ*.

were mistaken by Latreille for feelers, while the under lip forms a tube.



Parts of the mouth in a gnat magnified and separated.—*a, b*, the under lip and upper jaws carrying the jointed feelers; *f, c, d*, the under jaws and tonguelet; *e*, the under lip.

The parts are nearly of the same form as those of the gnat, in the common flea; and in gad-flies the arrangement is similar, though the parts are much stouter and more formidable.

In bugs, the under lip forms a long sheath; the edges, bending downwards, are shaped into a hollow canal, which receives the two pairs of jaws, in this case formed like long bristles, the two middle ones probably acting as piercers, while the other two probably assist in sucking.

MEMBERS OF THE CORSELET.

The members which belong to the corselet are appropriated to locomotion, and are of two sorts. One sort are jointed to the under pieces of each of the three rings of the corselet; these are the legs¹; another

(1) In Latin, *Pedes*.

sort are jointed into the upper pieces of the mid corselet and the hind corselet,—never to the fore corselet ; these are the wings¹.

Here the law of proportion is observed to regulate, as it always does, the relative size of the members, and the pieces on which they are jointed. When the legs, or a pair of them, accordingly, are very short, the corresponding pieces of the corselet are very small ; and when the wings are small, or wanting, the corresponding pieces in the corselet are small or undivided.

THE LEGS AND FEET IN INSECTS.

Insects have exactly six legs, though the first pair or the fore legs are in some butterflies so small as scarcely to be detected. Millepedes, centipedes, and wood-lice, which have more than six legs, are not strictly insects ; nor are spiders and mites, which have eight legs.

The fore legs, or first pair, are jointed into the jointings of the breast-plate with the under-flanks in the fore corselet, and are always directed forwards. The mid legs, or second pair, usually shorter than the others, are jointed with the same pieces in the mid corselet. The hind legs, or third pair, are jointed into the same parts of the hind corselet, are usually much longer than the first pair, and always directed backwards.

Parts of the Legs and Feet.

Each leg may be considered as made up of four principal pieces—the haunch, the thigh, the shank, and the foot—enclosed in a horny or membranous skin, containing the necessary muscles for moving the joints.

(1) In Latin, *Alæ*.

The haunch ¹ is various in form, being short and small in most beetles, while it is large in wasps and grasshoppers. In lady-birds it is round; in cockroaches it is flat. It is sometimes made up of three pieces, the two ² next the corselet working in the socket ³; and the third ⁴ jointed to the thigh, but apparently without independent motion.

The thigh ⁵ is the second principal piece, and is always comparatively long and generally bulged, but flat. It can only be moved backwards and forwards, and not sideways. In leaping insects, such as the green grasshopper, and in those which dig and burrow, such as the mole cricket, the thigh is always long, strong, and muscular.

The shank ⁶ is the third principal piece of the leg, and is generally very flat, and about the same length as the thigh, but more slender. In the hind legs of swimming insects, the shank is often fringed. Upon the upper end and sometimes on the middle, moveable spurs ⁷ are frequently jointed and fixed.

The foot ⁸ is the fourth principal piece, with which the leg ends. It is composed of five smaller pieces or joints ⁹ in a great number of species, but in some cases only from one to four, and in others, the number varies in the several pairs of legs on the same insect.

Systematists, following M. Latreille, have taken advantage of the variable number of these joints in the numerous species of beetles, to group them under several divisions.

The most remarkable part of the foot is the tip

(1) In Latin, *Coxa*.

(2) In Latin, *Trochantina* and *Rotula*.

(3) In Latin, *Acrotabulum*.

(4) In Latin, *Trochanter*.

(5) In Latin, *Femur*.

(6) In Latin, *Tibia*.

(7) In Latin, *Calcaria*.

(8) In Latin, *Tarsus*.

(9) In Latin, *Phalanges*.

piece termed the claw ¹, which is wanting in the fore-legs of some butterflies and other species. Each foot has usually two claws, but in some insects there is only one, and in others, there are four and even six on one foot. The claws are very various in form according to the uses intended, as we shall see presently.

The two winged flies with very long legs, popularly called jenny spinners, or daddy long legs, have the attachment of the haunch so slight, that they can at pleasure throw off the whole limb, as lizards sometimes do their tails, which I have often seen.



The leg of an insect. *a*, the haunch; *b*, the thigh; *c*, the shank, with a forked spur; *d*, the foot, with five joints.

Foot-combs and Foot-brushes of Insects.

Most people may have remarked, that flies are ever and anon brushing their feet upon one another to rub off the dirt, and equally assiduous in cleaning their eyes, head, and corselet with their fore legs, while they brush their wings with their hind legs. In the foot of the common blow-fly, there are two rounded combs, the inner surface of which is covered with down, forming a fine brush; and some crane flies have three such combs on each foot. Mr. Blackwall

(1) In Latin, *Unguis*.

denies that what are here termed 'combs exist, as they are represented by Bauer, on the edge of the membrane of the flies' foot, and says, the appearance of this is caused by minute short hairs. But though I should be more disposed to trust so celebrated a microscopical observer as Bauer, yet it is of little moment to the office of a comb, which I have here ascribed to the foot, whether it is hairs or serratures. I have also observed the water measurer clean its feet, though not by crossing them like the fly, but merely by wiping them upon one another.

In the case of the large green locust¹, the structure of the instrument is more obvious; and it being important to keep the soles of its feet clean, to enable it to walk up glass or other smooth substances, it is not contented, like the fly, with brushing them one on another, but will, like a spider, as first recorded by me, go over the foot with its jaws, mumbling off every bit of dust, a process which Swammerdam mistook for chewing the cud.

Muscles of the Legs and Feet.

Beginning with the haunch, Straus-Dürckheim describes in the cockchafer four flexor muscles in the fore leg, three in the mid leg, and five in the hind leg, without mentioning two others attached to the latter, which he supposes rather to be pertractors of the abdomen. Opposed to these are the extensor muscles of the haunch, one strong one in the fore leg; a short one and a long one in the mid leg; and three in the hind leg. Besides these, there are a flexor and an extensor to that which I have called the third piece of the haunch.

The thigh, in each of the three pairs of legs, has

(1) In Latin, *Acrida viridissima*.

only a very feeble abductor muscle, and as it is but very slightly moveable there is no antagonist adductor.

The shank has a strong flexor muscle, and a more feeble extensor in each of the three pairs of legs.

The foot, in each of the three pairs of legs, has an extensor and a flexor in each of the joints. The last joint includes a flexor and extensor for moving the claws.



Muscles of the right fore leg and foot in the cockchafer magnified eight times. Fig. 1, *a, b, c*, extensor muscles of the trochanter cut across; *d*, flexor also cut; *e*, the trochanter; *f*, extensor of the leg; *g*, flexor of the leg; *h*, extensor of the foot; *i, l, l*, the flexor of the foot, the tendon being the long white line in the centre; *k*, the leg; *m*, extensor of the claws; *n*, the flexor of the claws; *o*, the foot; *p*, the claws. Fig. 2. Hauch of the same laid open; *a*, the third flexor muscle; *b, c, d*, extensors of the trochanter; *e*, the flexor of the trochanter; *f*, the rotula; *g*, the trochanter. Fig. 3. Thigh of the same; *a*, portion of the extensor of the leg; *b, c*, the two side parts of the flexor.

Standing¹ of Insects.

The centre of pressure² in insects being usually either in the base of the abdomen or in the hind coxal joint, the three pairs of legs are of course inserted in advance of it; and hence, did they stand perpendicular to the body like the legs of quadrupeds, it would be impossible for an insect to alight on its legs without being upset.

To obviate this, the legs extend to a considerable length from the body, and not only so, but the first pair project obliquely forward, and the third pair obliquely backward, while the second pair are, in some species nearly at right angles to the body, and in other species, project obliquely forward or obliquely backward, according as the position of the centre of pressure requires for the stability of the animal. In two winged flies, for example, the mid pair of legs projects obliquely forwards, as may be seen in the figure.



Figure of a Crane Fly³ to show the position of the feet, as extending the base within which falls the line of direction of the centre of pressure.

(1) In Latin, *Statio*.

(2) See ALPHABET OF PHYSICS, page 88.

(3) In Latin, *Tipula oleracea*.

The singular pendulum-like motion, exhibited by some species of long legged flies while their feet are quite stationary, may be referred perhaps to the same cause as the similar motions of caged animals in a menagerie. When they are observed to perk up their hind legs in the air while they are standing on their fore legs—a common practice among gnats, it is probably to ascertain, from the agitations of the air, the approach of danger, as they are extremely timid.

Creeping, Walking, and Running of Insects.

The peculiarities of movement, in walking and running, exhibit an endless variety; for while some move straight onward, others move sideways, backwards, or even in circles and half-circles, while others walk on their back, others again suspended from a ceiling

Some insects, such as the oil beetle and the bloody-nosed beetle, walk very slowly, lumbering along at a pace which seems almost to indicate exhaustion and faintness, similar to that which affects bees during autumn, when they may often be seen crawling lifelessly upon flowers, as if so overgorged with the honey they had extracted, as to be incapable of flying. Some of the plant bugs both walk very slowly and by awkward jerks, having more the appearance of a bit of straw caught in a spider's web and waving irregularly in the air, than a living insect.

Many beetles run with great velocity, and dart off into holes and corners before a collector can catch them; but it is remarkable that those insects seldom run far without making a full pause to reconnoitre their position, as a deer may be seen to arch his neck from behind a tree to examine a stranger, and, after tripping off to some distance, turn round again to take another peep at the intruder.

Several of the midges, instead of walking straight forwards, walk obliquely backwards, and often at right angles to the line of their own bodies; while most insects can, when it is necessary, walk directly backwards with almost as much facility as forwards.

The bat louse, as described by Colonel Montagu, has its legs on its back, and, consequently must walk, which it does with miraculous agility, having its belly upwards, in the same way as several insects swim.

Suspensory Standing and Walking of Insects.

It is not a little curious to see the common house fly walk with ease up a perpendicular polished glass, and even remain suspended from the ceiling, with its back downwards. Dr. Power, in 1664, attributed this to a viscous matter squeezed from sponges in the flies' feet; Dr. Hook, in 1665, to minute particles and smoke on the glass, enabling the flies to hold thereby; and Dr. Derham, in 1711—12, to suckers which exhausted the air, and consequently produced pressure from the atmosphere¹.

The last opinion was adopted by White of Selborne, and more recently illustrated by Sir E. Home and M. Bauer, who have described and figured the suckers in the foot of the fly, supposed to act in the same way as the leather sucker with which boys drag stones.

Mr. Blackwall endeavoured to put this plausible opinion to the test of experiment, and found that when the flies were by any means enfeebled, they could not walk on glass which was highly polished, but succeeded on what was slightly soiled or defective in polish. What was more extraordinary, the house fly was found to traverse the exhausted receiver of

(1) See ALPHABET OF PHYSICS, page 81.

an air pump in all directions, so long as it remained vigorous. He thinks, therefore, that as the best polished glass is not free from flaws, and as what has been called the suckers in a fly's foot, are fringed with hairs, these lay hold of the inequalities in the glass.

I am of opinion, however, that though Mr. Blackwall has partly shaken the explanation of Derham and Sir E. Home, he has not at all established his own. It would never apply, I am confident, to the green locust¹, which I have often seen walk with ease up a pane of glass.

Swimming of Insects.

The bodies of most insects being lighter than water, it requires no effort to buoy themselves in swimming, and they only require to strike the water like other animals with their legs in order to swim. Those which dive easily probably do so by compressing the air vessels within their bodies. The fore feet are rarely used in swimming, inasmuch as they would tend to produce retrograde movements; the hind feet being almost exclusively used.

The greater number of water insects, however, rather walk than swim, and most people may have observed groups of them walking up against small streams, apparently more for the purpose of maintaining their place, than of making farther progress. Amongst these are the water measurer, several water bugs, as well as gnats, all of which have such slender feet, that I can only explain their not sinking in water on the same principle as that of a small needle floating when very dry and laid exactly level—which is buoyed up by a thin layer of air between it and the water.

(1) In Latin, *Acrida viridissima*.

The whirl-wig beetle, as I have elsewhere said, can scarcely be said either to walk, run, or swim,—for, as I have taken some pains to ascertain, it appears not to keep its “oary feet” plunged in the water as it flits about, but strikes smartly out, and suddenly folds them flat under its belly, on the same principle as a waterman on the Thames may be seen to give two or three quick pulls with his oars to put *way*, as he calls it, upon his wherry, and then, perking them up out of the water, lets it skim along while he rests motionless. Thus does the little whirl-wig glide along the water, as if by magic; for we cannot see its feet moving on account of the border of the wing-cases which overlap them; no more than we can discern the feet of a swan, from their dark colour, resembling that of the water, even when swimming about at a small distance from the shore.

Many insects walk at the bottom of water, in the same way as they do on land.

Hopping and Leaping of Insects.

Several winged insects, in the same way as many birds, instead of walking, leap or hop along somewhat in the manner of a kangaroo or jerboa. This motion, as M. Straus-Dürckheim remarks, is performed by the rapid unbending of the hind legs forcibly pressed for the purpose against the substance whence the leap is made.

Some species make use of their faculty of leaping to escape from their enemies, as well as for the purpose of changing places, while others use it to spring on their prey.

The most common and best known instance of a leaping insect is the flea, the extraordinary muscular power of whose thighs enable it, not only to perform

most astonishing leaps, but when harnessed to a small coach or brass cannon, seventy times its own weight, to draw like a horse, as I have witnessed. A flea will leap two hundred times its own length, but in this it is outdone by the cuckoo-spit froghopper, which can leap two-hundred and fifty times its own length, while a man could not leap any thing near this, which in his case would be three or four hundred yards.

The minute observation, by which such unexpected facts are discovered, has in all ages been a fertile source of ridicule for the wits, from the time when Aristophanes, in his *Clouds*, introduced Socrates measuring the leap of a flea, up to Peter Pindar's lampoon on Sir Joseph Banks and the emperor butterfly. To such flippant wit, we have merely to retort the question of the Abbé de la Pluche, "if the Deity thought insects worthy of His divine skill in forming them, ought we to consider them beneath our notice?"

THE WINGS IN INSECTS.

The wings in insects are four, sometimes two, in number, and are uniformly jointed upon the upper flanks, and the back plate of the corselet,—one pair uniformly on the mid corselet, and the other uniformly on the hind corselet.

The upper or fore wings¹, which may be also called the greater or first pair, are always jointed upon the two upper flanks, and the back plate on each side of the mid-corselet, and of course near the centre of pressure², being balanced on the one side by the head and fore corselet, and on the other by the abdomen and hind corselet.

(1) In Latin, *Alæ anteriores*.

(2) See ALPHABET OF PHYSICS, Page 88.

Parts of the Wings.

Somewhat like the human wrist, which is composed of a number of little bones, the wing joint in insects has small horny joint pieces³, to the number of seven in the first pair, of various size and figure, but all united by a membrane, and jointed on the one side with the wing, and on the other, with the back plate and upper flank.

These joint pieces move the wings by means of three muscles,—the first divided into two portions where it is attached to the inside of the corselet; but these unite into a single tendon, and are fixed into one of the joint pieces. When this muscle contracts, it lowers the base of the wing and consequently raises the tip.

A second muscle is also fixed within the corselet, and is attached to another of the joint pieces. This muscle lowers the inner edge of the wing and gives it a sort of pendulum-like motion.

A third muscle is similarly placed, and acts in concert with the two first.

The preceding description applies to the wings of bees and wasps; while in two-winged flies M. Chabrier describes two sets of muscles: one set placed lengthwise for lowering the wings, and another placed aslant and across these for raising the wings.

M. Straus-Dürckheim describes no fewer than thirteen pairs of muscles, as employed, directly or indirectly, in moving the second pair of wings in the cockchafer; amongst others, retractors, depressors, elevators, pertractors, extensors, relaxors, and flexors, independently of those to be afterwards noticed, which move the wing cases.

(3) In Latin, *Epidemæ*.



Interior of the corselet and muscles of the under wings in the cockchafer magnified eight times.—*a*, rotator of fore corselet; *b*, retractor of the mid back-plate; *c*, retractor of the wing; *d*, depressor of the mid back-plate; *e*, extensor of the wing case; *f*, flexor of the wing case; *g*, first flexor of the mid haunch; *h*, its short extensor; *i*, its long extensor; *k*, long head of extensor of the trochanter; *l*, second head of elevator of the wing; *m*, anterior extensor of the wing; *n*, the posterior extensor; *o*, retractor; *p*, relaxor of the wing; *q*, second flexor of the hind haunch; *r*, the fifth flexor; *s*, third extensor; *t*, part of long head of extensor of trochanter; *u*, short head of the same; *v*, flexor of the trochanter.

In general, the wings may be said to be composed of two membranes united together by means of horny lines, which are variously termed veins, nervures¹,

(1) In Latin, *Nervæ*.

and wing bones, though not quite correctly. I prefer the term rib¹.

Disposition of the Wing Ribs.

The wing rib is not to be considered as a bone, but as a horny and nearly solid tube enclosing air-pipes for the purpose of expansion. The number and disposition of these ribs, which may be reckoned about seven, vary greatly in different groups of insects.



a, wing of dragon fly; *b*, wing of bee; *c*, wing of a house fly.

In bees, wasps, two-winged flies, and butterflies, beginning at the upper edge of the wing, we find it formed by a strong rib which may be termed the fore-rib³, rendered remarkable in some butterflies for a hook at its base, which serves as a pulley for the tendon of a muscle.

At some distance, often about the middle of the wing, and somewhat parallel to the fore rib, runs another also usually strong, sometimes the strongest of all the ribs,—which may be termed the mid rib⁴,

(1) In Latin, *Costa*.

(2) In Latin, *Tracheæ*.

(3) In Latin, *Costa anterior* or *Radius*.

(4) In Latin, *Costa media* or *Cubitus*.

for though not quite in the middle of the disc of the wing, it is for the most part nearly so at its base.

In many insects, particularly bees, butterflies, moths, and dragon-flies, the fore and mid rib join near the upper edge, at some distance from the tip, forming a small opaque horny plate, probably serving as a reservoir for air or fluid, which may be termed the rib-spot¹.

From near the base in the mid rib, there is often given off a branch which runs between it and the fore rib, sometimes, as in butterflies, so large, that it may be mistaken for the mid rib itself. It may be termed the mid rib branch². It is wanting in bees.

On the lower side of the mid rib, several branchlets are given off, which go to unite with other branchlets. These branchlets may, if necessary, be numbered 1, 2, 3, &c., or lettered *a*, *b*, *c*, &c.

The next leading rib in the wing may be termed the inner rib³, and is usually divided near the base into two, sometimes three or four, branches. In butterflies, the division takes place about the middle of the wing.

Near the under edge of the wing is another rib, which may be termed the lower rib⁴, between which and the edge, are in some groups one or more small ribs⁵.

These several ribs and their branches, (in some groups so numerous as to give the whole wing the appearance of lace or net work,) by inclosing portions of the wing, form what may be termed areas⁶, and as it is of some use to attend to these, in describing the

(1) In Latin, *Stigma*.

(2) In Latin, *Costa mediantina*.

(3) In Latin, *Costa interna*.

(4) In Latin, *Costa inferior*.

(5) In Latin, *Costulae marginales*.

(6) In Latin, *Areae*.

wings, I shall point out the principal areas, premising that a wing may be said to have a base ¹, where it is joined to the corselet; an upper edge ² running from the base to the fore tip ³, an outer edge ⁴ running from the fore to the hind tip ⁵, and a lower edge ⁶ from the hind tip to the base.

What may be termed the upper area ⁷ is the space inclosed between the mid rib with its branches and the upper edge.

The space inclosed between the mid rib and the inner rib, with its branches, may be termed the mid area. In butterflies, this space ^b is somewhat oblong and triangular, without any cross ribs.

The space inclosed between the mid rib and the lower rib, or between the lower rib and the under edge, may be termed the lower area ⁹.

The irregular space, occupying the outer edge; all the wing, indeed, not included in the three other areas, may be termed the outer area ¹⁰.

The several areas are further divided into smaller areas, or, as they may well be called, meshes ¹¹, which form beautiful lace-work in the dragon flies and others of the same group.

According as they are in the base, the middle, or the outer edge, these may be termed base meshes ¹², mid meshes ¹³, and outer meshes ¹⁴.

Markings of the Wings.

The other parts of a wing necessary to be attended

(1) In Latin, *Basis*.

(2) In Latin, *Margo anterior*.

(3) In Latin, *Apex anterior*.

(4) In Latin, *Margo exterior*.

(5) In Latin, *Apex inferior*.

(6) In Latin, *Margo inferior*.

(7) In Latin, *Area superior*.

(8) In Latin, *Area media*.

(9) In Latin, *Area inferior*.

(10) In Latin, *Area exterior*.

(11) In Latin, *Reticulæ*.

(12) In Latin, *Cellulæ basilares*.

(13) In Latin, *Cellulæ mediæ*.

(14) In Latin, *Cellulæ exteriores*.

to in descriptions, are chiefly coloured markings, particularly on the wings of butterflies, such as a streak ¹, a band ², a spot ³, an eyelet ⁴, a sprinkling ⁵, &c. The outer and lower edges, are, in butterflies, moths, and some other insects, furnished with fringes ⁶, and sometimes with a tail ⁷.



Markings and Edgings of the wings of butterflies and moths.

Texture of the Wings.

With respect to texture, the first pair of wings in beetles is so horny and stiff, that they have no close resemblance to what are called wings in other insects. They are indeed less adapted to flight than to cover the second pair in a state of repose, and to balance the body, perhaps, by their weight in flying. Sometimes indeed these horny wings, usually called wing cases ⁸,

(1) In Latin, *Striga*.

(3) In Latin, *Macula*.

(5) In Latin, *Irroratio*.

(7) In Latin, *Cauda*.

(2) In Latin, *Fascia*.

(4) In Latin, *Ocellus*.


(6) In Latin, *Ciliæ*.

(8) In Latin, *Elytra*.

cannot be separated on the back, in which case the second pair of wings is either very small or altogether wanting. The term wing cases is quite appropriate, so far as function is considered; but it is important to know that these wing cases in beetles occupy the same place, and are jointed in the same manner, as the first pair of wings in bees.

M. Straus-Dürckheim describes the wing cases of the cockchafer as being jointed by means of four little joint pieces, and numbered first, second, third, and fourth. Two of these are moved by appropriate muscles.

The wing cases are themselves put in motion directly by a pair of extensor muscles, a pair of adductors, and a pair of flexors; indirectly by the retractors and depressors of the mid back plate of the corselet.



The first or inner layer of muscles of the corselet and abdomen in the male cockchafer. 1, 2, 3, 4, 5, 6, 7, 8, the rings of the abdomen; *a*, the elevator of the head; *b*, the depressor; *c*, the rotator; *d*, the lateral flexor; *e*, the retractor of the jugular; *f*, the oblique elevator; *g*, the strait elevator; *h*, the upper retractor of the fore-corselet; *i*, the lower retractor; *k*, the elevator; *l*, the rotator; *m*, the first flexor of the haunch; *n*, the second flexor; *o*, the fourth flexor; *p*, the extensor of the trochanter; *q*, the retractor; *r*, the extensor of the wing-case; *s*, the short extensor of the middle haunch; *t*, the pertractor; *u*, the lateral flexor; *v*, the depressor of the back-plate; *w*, the depressor of the midriff; *x*, the depressor of the wing; *y*, the elevator; *z*, the second flexor of the hind haunch.

The leathery wings of locusts, crickets, and grasshoppers, and the half leathery wings of tree bugs, and some other insects, are also similar in their relative situations to the first pair of wings in bees.

Peculiarities in the Wings.

In the base of the first pair of wings, there are found in the common fly, some beetles, and other insects, a pair of small scaly members, dependent on the larger wings, and appropriately termed winglets¹. It is a mistake to consider these apart from the wings, for they are always jointed into the back plate of the mid corselet, never into the hind corselet, and united at the base to the wing; and often the separation of one of them from the wing cannot be traced. It is sometimes double, that is, two to each wing, like a bivalve shell. It does not as has been supposed produce the buzzing of flies.

The second pair of wings² are always, like the first pair, jointed with the back plate and upper flanks of the hind corselet. They are in most respects similar to the first pair, as in the joint pieces, which are only six instead of seven, the ribs, the areas, and the parts of the circumference.

It is worthy of remark that, except in dragon flies, earwigs, some beetles, and a few other insects, they are commonly much smaller than the first pair, but in all cases following the law of proportional relation to the parts of the corselet upon which they are jointed.

In the common fly, and others similarly constructed, there arises exactly from the same part of the hind corselet as the second pair of wings, a pair of small members, shaped somewhat like a drum-stick, and protected above by the winglet. These have been

(1) In Latin, *Alulae*.

(2) In Latin, *Alae posteriores*.

called poisers¹, or balancers, and are supposed by some to aid in balancing the body, by others to produce buzzing, by beating on the winglets, neither of which opinions is proved. The law of proportion assimilates them, as M. Audouin thinks, to the second pair of wings, which he proves by the fact of the poisers only occurring in insects having no second pair of wings.

In moths, the second pair of wings are each furnished near the base with a sort of slender, but stiff, horny hook, somewhat curved, which is fixed into a projection of the first pair by way of bridle².

In wasps and bees the second pair are bridled to the first pair by minute hooks, in form of an S along the upper edge.

The wings of some insects are clothed with hair, and others, as butterflies and moths, with a sort of feathery scales³, which appear in the microscope of very various forms. These scaled feathers are placed over each other like the tiles of a house, are the coloured part of the wing, and easily come off on being touched in the form of dust.



The feathery scales from the wings of butterflies and moths, magnified to shew their various forms,—from Reaumur.

(1) In Latin, *Halteres*.

(2) In Latin *Frænum alæ*.

(3) In Latin, *Plumulæ*.

Flying of Insects.

When a pigeon or a pheasant rises to fly, it may be observed, first to fill its body with air by taking several deep inspirations, at the same time often rising on tiptoe, hopping a short space, and puffing out and balancing its body to feel whether it be light enough to rise. In a similar way, many butterflies after alighting may be seen to flirt their wings as a lady does her fan, for the purpose, as I conjecture, of throwing air into their bodies previous to flight. I am aware that M. Chabrier, who wrote a quarto volume on the flight of insects, has retracted this notion respecting the air; but I am borne out in adopting it by the still higher authority of M. Straus-Dürckheim.

The motion of flight in insects is produced solely by alternately raising and lowering the wings, and never by extending and retracting them, as may be observed most easily in the dragon fly, whose wings are always extended. In beetles, the position of the wings and of the wing cases, causes the body, during flight, to be oblique rather than horizontal. Crows fly in the same way as insects; but woodpeckers fly by giving alternately three or four rapid jerks with their wings, and then rest.

A pretty group of two winged flies, somewhat resembling wasps in colour and size, may frequently be remarked in lanes and woods, hovering on the wing for a considerable time, without shifting a hair's breadth from their place, though the motion of their wings is all the while so rapid, as to be almost imperceptible, similar to that of some moths whilst sipping on the wing the honey of flowers. The vibratory motion of the wings, in these cases, is very different from that of flying, as may be proved by alarming them, when

they will be observed to change the vibratory for a verberatory motion, and make their escape. The proceedings of the kestrel hawk or windhover are exactly alike.

The flight of some insects is very slow, as in the instance of the wood white butterfly; and very rapid in others, as in the dragon fly. I observed some water insects this summer, (1833,) skimming the brook at the bottom of my garden so rapidly, as to produce the appearance of numerous grey lines along the water, in the same way as a piece of burning charcoal whirled rapidly round, produces the appearance of a circle of fire.

The gilded summer flies, which mix and weave
 Their sports together in the solar beam,
 Or, in the gloom of twilight hum their joy;

have been supposed to congregate for various reasons, none of which seem tenable, except that they are produced in numbers at the same time in the same place. It has been remarked that the gnats, which dance in chorus, are uniformly males; and I am very certain that it is only male butterflies which may be seen sporting together in the air in mock skirmishes.

These butterfly skirmishings do not appear to me to be prompted, as Mr. Knapp seems to suppose, by testiness nor jealousy, but by the spirit of gaiety and frolic;—the buoyant feelings arising from the air expanded in their wings and bodies by the warm sunshine causing the living principle to increase, even to exuberance. Were these, indeed, actual combats among the males, like those which take place among game-cocks or ruffs, nature would probably have furnished them with weapons suited to their warfare: but butterflies have neither spurs, claws, nor sharp bills, wherewith to assail an enemy; and though they

might flap one another with their wings, till their tiny feathers flew about like a snow-shower, yet I never have observed them do so.


MEMBERS OF THE ABDOMEN.

THE only members in this portion of an insect's body are situated on the two last rings, and are very various in form and use. Insects, having no spine like animals furnished with bones, have no tail, at least similar in structure to that of the dog or the horse; but in many moths there is a hairy brush, which is termed the tail¹, on the last ring. Its use is not apparent in the males, while in the females the hair is plucked out to cover their eggs, by an instrument similiar to a pair of tweezers, also placed on the last ring of the abdomen. In the scorpion fly the tail is jointed.

In earwigs there is a forked member² on the last ring, the blades of which are moveable, and which are said to be used for folding up (rather, I should think, for unfolding) the wings, which are for the most part concealed under the short wing cases. A somewhat similiar member, but the blades crossing each other as scissors, is found in the male snake fly. In the rove beetles are long, narrow, stiff, members; and in the dragon-flies, leaf-like members on the last ring, absurdly called a sting, but whose use is not well ascertained, any more than the use of the long bristles jointed into the last ring of the day fly, and shorter ones in the cockroach, and some crickets and grasshoppers.

(1) In Latin, *Cauda*.

(2) In Latin, *Forceps*.



(Back and front views of the Sugar-louse magnified.)

The antennæ (*f, f,*) are short, cylindrical, tapering from the base to the tip, and beset with bristles at the joints. The head is covered on the upper part with a round plate, and the back with twelve plates of a square form: the under part of the body, reckoning from the head, has only eleven such plates. The feet are connected with the three anterior divisions of the body, and consist of five joints, of which the two first are very broad in the middle, and the last very thin and roundish. On the hinder extremity of the body, are two short, thin, bristles (*palpen, q, q,*) beyond which, upon each side, are two very long ones (*r, r,*) and beyond these four others (*m, a,*) shorter, while the last ring of the body ends in a single pointed bristle (*schwanzspitze, t,*) similar to the rest, all of

them having precisely the same structure as the antennæ. The three longest bristles take their origin from the last back plate, under which is the vent; but the short bristles originate from the last belly-plate (*bauchschuppe*), between the two halves of which are the external organs of reproduction.

In other instances, there are long members in the last ring, somewhat similiar to a brad-awl, and used for the purpose of boring holes to deposit the eggs. These are peculiar to females, and are termed ovipositors¹. They are sometimes hollowed into a tube for the egg to pass along, and sometimes solid.

In bees, wasps, and some other insects, there is a weapon used for attack and defence, termed the sting², which can be drawn within the abdomen when it is not used. It is composed of a sheath, darts³ barbed at the point, and a poison bag at the base for poisoning the wound which it makes.

In the grub of the glow-worm, I discovered a singular instrument on the last ring, composed of gristle-like rays, in form of a funnel, covered with a clammy adhesive substance, and capable of being extended and contracted. The grub employs it to clean every part of the body.



a, The last ring of the abdomen of a bee opened, shewing the sting in its sheath; *b*, the sting of a bee magnified to show the barbed darts; *c*, cleaning instrument of the grub of the glow worm open; *d*, the same shut.

(1) In Latin, *Ovipositores*.

(2) In Latin, *Aculeus*.

(3) In Latin, *Spicula*.

INTERNAL ORGANS OF INSECTS.

HAVING thus taken a brief, but, I trust, an intelligible survey of the outside of the body in insects, the student may be supposed to be partly prepared to examine the various organs within the body. It will be convenient, for this purpose, to begin with the organs employed in the digestion of the food, and then pass on to those employed in breathing, sensation, and reproduction. The greater number of the muscles have already been noticed under the several motions which they produce.

ORGANS OF DIGESTION.

IT is remarkable that the length of the organs of digestion in insects, measuring from the mouth to the vent, is, as in the larger animals, proportioned to the sort of food. Vegetable food, being more crude, or not so like the properties of the animal body, requires more preparation to turn it into nourishment; and hence insects, and other animals¹ which feed on vegetables, have their organs of digestion of great length, much longer, indeed, than the body, in which they wind in many folds. Animal food requiring less preparation, the insects which feed on it have their

(1) See ALPHABET OF ZOOLOGY, page 72.

organs of digestion short, and of the same length as the body. In all cases they consist of three layers, the outer membranous, the middle muscular, and the inner mucous.

When the food has been taken into the mouth and bruised, or chewed by the jaws when those are moveable, or sucked up when they form a sucking tube, it passes on to the *haus*¹, then to the entrance² of the gullet, and then to the stomach and intestines, as in the larger animals. It is worth remarking, that in most instances the structure of the stomach and intestines is nicely conformable to the structure of the mouth.

In man, the food is mixed, during the process of chewing, with a peculiar fluid supplied from several glands or fountains situated near the mouth. In insects, similar fountains³ have been described by Ramdohr, Leon Dufour, and others. They are most obvious in sucking insects, and when a fly cannot suck a bit of dry sugar, it has been observed to moisten it with this fluid.

The organs⁴ which furnish the silk, spun by the silk-worm and other caterpillars, are similarly situated with the preceding, and perhaps are the same organs.


The most complicated organs of digestion, (found, of course, in insects feeding on vegetable matter,) may be described under six divisions, similar to those of birds, namely, the gullet, the crop, the gizzard, the stomach, the intestines, and the vent.

(1) In Latin, *Fauces*.

(2) In Latin, *Pharynx*.

(3) In Latin, *Sialistaria*.

(4) In Latin, *Sericteria*.



The organs of digestion in two different beetles. Fig. 1, a, garden beetle (*Carabus*). Fig. 2, a churchyard beetle (*Blaps*). a, the jaws and feelers, b, the head; c, c, the saliva vessels; d, the gullet, very short in Fig. 1; e, the crop, wanting in Fig. 1; f, the gizzard, wanting in Fig. 1; g, the stomach, large and convoluted in Fig. 1; h, h, h, the bile vessels long, and numerous; i, the small intestine; k, the blind gut; l, the vent; m, m, the excrementary vessels.

The gullet ¹ varies much in length, being sometimes very short, and sometimes reaching to the abdomen, or even within it, but it is more generally the length of the fore corselet within which it is lodged. At its upper end, it is surrounded by a nervous ring, from which two branches go off and unite at the lower part of the body. It ends in the stomach ², which is sometimes simple, and sometimes composed of three chambers.

The crop ³ or craw is a bulging out of the gullet into a sort of pouch, which, on the outside, can seldom be distinguished from the gizzard; but, on the inside, it is found destitute of horny projections, and its texture is more membranous than muscular. When large, it is often seen with folds or plaits. It constitutes what is called the honey bag of the bee, and it is in it that insects have the fluids, often offensive, which they discharge when caught, as is observed in many beetles. In butterflies and other sucking insects, it is placed on one side of the gullet, and not in the line of the stomach. Its contents have, therefore, to be returned into the mouth before they can reach the stomach. The crop is not found at all in many instances.

The gizzard ⁴ succeeds the crop, and is more muscular in structure, and furnished on the inside with moveable horny projections, most probably employed in bruising the food. These horny pieces are of various figures, and placed in various directions, sometimes like a brush, sometimes like a comb; and just above the entrance of the stomach they nearly close the pas-

(1) In Latin, *Œsophagus*. (2) In Latin, *Ventriculus*.

(3) In Latin, *Inglucies*; in German, *Speis sack*.

(4) In Latin, *Echinus*.

sage, forming a sort of valve, which will only permit minute portions of the food to pass. Swammerdam and Cuvier are mistaken in thinking insects which have a gizzard, such as grasshoppers, chew the cud. The chewing motion which deceived them, I have found to be the process of cleaning the feet and the ears. The gizzard is not found in all insects.

In the sugar louse the gizzard is furnished on the inside with six teeth, some hooked and calculated to seize the food, others formed so as to bite and chew it.



Gizzard of sugar louse laid open to shew the teeth. 1 and 4, biting teeth; 2 and 6, chewing teeth; 3 and 5, seizing teeth.

There are, according to Treviranus, similar stomach teeth of various structure in the oriental cock roach.

The third chamber, or digestive stomach¹, is composed of thin, soft, extensible, membranes, usually cylindrical in form, and sometimes forked, the entrance² being at one of the forks. One remarkable circumstance is, that, in many insects, the outer surface is covered with a number of teat-like points, similar to the finger of a glove, containing fluid which they discharge into the stomach. These may be termed vessels³. They vary much in size and number, and are not found in all insects. Different opinions are held respecting them by Cuvier, Marcel de Serres,

(1) In Latin, *Faliscus*.

(2) In Latin, *Cardia*.

(3) In Latin, *Villi*.

and Leon Dufour. This is the structure also when there is only one stomach.

In all insects, we believe, there are vessels called bile vessels¹, consisting of several membranous tubes, filled with a peculiar fluid, bitter, and usually brown or yellow, but sometimes limpid, supposed to be similar to bile, though we find nothing like the liver for preparing this bile, which is probably, therefore, as M. Gaëde thinks, prepared in the vessels themselves. These vessels float in many convolutions in the abdomen, one end being sometimes free and the other fixed, and sometimes both ends fixed, and giving rise to a sort of arch or curve. Sometimes these are inserted into the stomach near its outlet; in others, one end goes into the stomach, and the other into the blind gut. Their number varies from two, which are found in the rose-chafer, to four, found in beetles and common flies; to six, found in butterflies, and to even one hundred and fifty (probably mere branches of two fundamental ones), found in bees, wasps, and dragon-flies.

It is in the stomach that the food is converted, by means of the digestive fluid, from the gastric vessels and the bile, into a pulpy mass, called chyle, if we may follow the analogy of other animals.

The outlet² of the stomach is in insects furnished with a valve to prevent the too rapid passage of the chyle into the intestines.

The intestines³ form an extended portion of the organs of digestion, which may be divided into four parts, the chyle gut, the small gut, the blind gut, and the vent gut.

(1) In Latin, *Vesiculi biliarii*. (2) In Latin, *Pylorus*.

(3) In Latin, *Intestina*.

The chyle gut¹, which is always found in large animals, is seldom, in insects, different from the small gut. When it is distinguishable, as in the glow-worm, it is very smooth. It receives the chyle from the stomach.

The small gut² is usually straight, smooth, and of equal size through its whole length, though there are bulgings in some species; and, for the most part, it has many convolutions. The chyle, in passing along, has its nutritive portions taken up by the inner membrane of this intestine, through which it passes into the red blood, which is not found in insects.

The blind gut³ consists usually of an egg-shaped cavity, formed by the bulging out of the lower end of the small intestine. It is often covered with plaits or bands; sometimes the bile vessels open into it, and it always contains the crude parts of the chyle rejected by the small gut as unfit for nourishment. Insects have no gut precisely similar to the colon of other animals.

The vent gut⁴ is very muscular, and usually short. It ends in the vent⁵, through which the crude parts⁶ of the chyle, collected in the blind gut, are thrown out of the body.

The singular discharges of offensive matter, such as the poison of the bee, and the vapour of the bombardier beetle, are prepared near the vent gut by a particular apparatus, and stored up in a sort of bladder, from which they are discharged.

In the abdomen of a certain class of bees in the common hive, called wax workers, are cells between the rings, in which wax appears to be prepared by

(1) In Latin, *Duodenum*. (2) In Latin, *Intestinum tenue*.
 (3) In Latin, *Cæcum*. (4) In Latin, *Rectum*.
 (5) In Latin, *Anus*. (6) In Latin, *Excrementa, or, Fæces*.

secretion from the food within, and not collected, as is supposed, directly from flowers; as the pollen is well known to be upon the thighs for the purposes of food. This opinion, however, which I have not myself verified by observation, is contrary to that popularly held. It was first started by Hornbostel, a clergyman at Hamburgh, in 1744, and republished as his own discovery, by Reim, in 1769. Mr. John Hunter, evidently without being aware of these, published it as his own discovery in 1792; and Huber, assisted by the clever daughter of Professor Jurine, made experiments and dissections, all confirmatory of the same view. Recently G. R. Treviranus, one of the best living experimental physiologists, has repeated the investigations, and has come to the same conclusion. I think these high authorities must outweigh that of Mr. Huish, who decides that they are all wrong, and that the popular notion is right.

The nutritive part of the chyle, which is (if the term may be used) filtered through the sides of the small gut, is not then received into any vessel, as has already been said, but spreads about through the interior of the body, taking the form of an irregular mass of soft pulpy fat, greenish or whitish in colour, which surrounds the organs of digestion, and fills up every vacant place in the body, particularly in caterpillars, of which it forms a very large proportion of the whole bulk.

The more fluid portions appear to be taken up through the membranes of the several organs, probably in a similar way to that by which they previously passed through the membranes of the small intestine. On this subject, however, we are still very much in the dark: though it is certain the fat is employed for the purposes of nutrition; for it is always stored up in great quantity before an insect passes

into the state of chrysalis, when it ceases to eat, and often remains torpid for many months; and is also found similarly stored up in the female before the eggs are laid, but after this disappears.

The nutritive matter, whether it be fat or fluid, not being carried through the body in arteries and veins as the blood is in other animals, to nourish the several parts, lies around and upon the parts to be nourished, which absorb the peculiar portions they require, rejecting the rest; and this may be required by other parts for a different purpose.

The more solid portions may be required by the muscles, the layers of the intestines, the horny skin, and perhaps by the nerves; while the more fluid portion may be taken up by the gastric vessels, the bile vessels, and the peculiar vessel, which is the only organ found in insects in the least resembling a heart.

ORGANS OF CIRCULATION.

THE organ or vessel which was called a heart by Malpighi, Swammerdam, and the older naturalists, is now commonly called the dorsal vessel¹ by modern writers; but I shall use the old term heart, which, besides being again introduced by Meckel, Herold, and Straus-Dürckheim, is less repulsive to a beginner; and I shall endeavour to describe it so as to prevent misconception.

The chief resemblance which the heart in insects bears to that of other animals, is its containing a fluid, and its regular beating, as may readily be observed in smooth caterpillars, in which it is of large size. Lyonnet counted twenty to one hundred beats in the minute in the goat caterpillar, caused, it has been said, by the alternate contraction of a number

(1) In Latin, *Vesicula dorsalis*.

of muscles ranged along its sides. It differs most essentially from the heart in other animals, in having no visible inlet or outlet in the form of veins or arteries. Consequently, there is not, and cannot be, any real or direct circulation of blood in insects. though a claim to the discovery of such a circulation was made, in 1824, upon very slight and vague grounds, by Professor Carus, of Dresden, and too hastily admitted by Mr. Spence. "People," says Goethe, "are very busy in repeating what has been discovered by others." In 1833, a similar claim was put forward for Mr. J. Bowerbank, and was loudly alleged as a proof of the inaccuracy of the first edition of this Alphabet; but the author has not stated a single fact not previously well known; while he expressly and correctly admits that "the blood does not appear to be confined within any specific vessels," before it passes into the heart. Dr. Carus uses nearly the same terms, and farther admits that there is only a circulation in the grub, not in the adult insect.

Without detailing minutely the vague and loose statements of Baker, of Lesser, of Comparetti in 1800, and of Goring and Pritchard in 1829; or the careful experiments of Lyonnet, Baron Cuvier, and Marcel de Serres, I shall content myself with quoting, in corroboration of what I have always maintained, the opinion of M. Latreille, in 1829, when he was aware of the claims of Dr. Carus. "The presence," he says, "of air pipes (*trachées*) excludes all complete circulation, that is to say, the distribution of the blood to different parts, and its return from the organs of respiration to the heart." This fully ascertained fact must render any claim to the discovery of an insect circulation absurd¹.


(1) See FIELD NATURALIST'S MAGAZINE, for May, 1833, p. 226.

The heart lies along the whole extent of the back, from the head to the vent, immediately under the skin and muscles. It is in form of a cylinder contracted at the two ends, but usually narrower towards the head, and wider towards the vent.

It is composed of two membranes; an outer of cellular texture interlaced with several air pipes¹, and an inner of muscular texture.

Within it is filled with a fluid, transparent, coagulable, readily drying, and, when dry, having the look of gum, its colour being sometimes strong and sometimes greenish, orange yellow, or dull brown. Without, the masses of fatty matter, which sometimes surround it in considerable quantity, are of the same tint as the fluid within, and it may be that the fluid derives its colour from them, though M. Audouin seems to think the reverse is the case.

Meckel and Herold, considering the organ a heart, think that its beatings, or alternate contraction and dilatation, which affect its whole extent, are for the purpose of agitating, not of circulating, the fluid it contains, as they admit no outlet or inlet by means of vessels for that purpose. Herold thinks that the dilatation, or diastole, is produced by the triangular muscles which attach it to the back; while the contraction, or systole, is produced by the muscular fibres of the inner membrane.



The heart of a carnivorous grasshopper, with its valves, chambers, and artery, from M. Audouin.

(1) In Latin, *Tracheæ*.

“The dorsal vessel,” says M. Straus-Dürckheim, “is in reality the heart of insects, being in them, as in higher animals, the mover of the blood, which, instead of being contained in blood vessels, is diffused through the cavity of the body. The heart occupies the whole of the back of the abdomen, and ends at the fore part of the organ in a single artery¹ without branches, which carries the blood into the head, where it is poured out, whence it flows back into the abdomen, in consequence solely of its accumulation in the head, in order to enter anew into the heart. To this alone is reduced the whole circulation in insects, which accordingly have but one artery without branches, and no veins.

“The wings of the heart are not muscular, as Herold pretends; they are simple fibrous ligaments for retaining the heart in its place.

“The heart is divided within into eight successive chambers, in the cockchafer, separated from each other by converging valves, which allow the blood to pass forwards from one to another as far as the artery that conducts it into the head, but oppose its movement backwards. Each chamber is furnished at its fore part on the sides with two openings in form of cross chinks, which communicate with the cavity of the abdomen, and through these the blood contained in the latter can enter into the heart. Each of these openings has within it a little valve, in the form of a half circle, which shuts up the passage during every contraction.

“From this brief detail, it may be conceived, that when the chamber nearest the vent is dilated, the blood in the cavity of the abdomen will enter through the

(1) This was anticipated in part by Lyonnet.

two openings¹ above described. When this chamber contracts, the blood which it contains, not being able to return into the cavity of the abdomen, shuts the valve², and passes into the second chamber. This dilates for the reception of the blood, and at the same time it receives a certain quantity of blood by its proper openings. Then, by the contraction of this second chamber, the blood passes in the same manner into the third, which receives it equally by the side openings. In this way the blood is propelled from one chamber to another, till it reaches the artery; and it is the successive contractions of the several chambers of the heart which are observable through the skin in the backs of caterpillars."



The heart of the cockchafer, with its valves, chambers, and artery, magnified eight times, from M. Straus Dürckheim.

This is both very rational, intelligible, and supported by minute and careful observation. It at once destroys numerous fancies on the subject, proposed by Baron Cuvier, Marcel de Serres, and others.

The blood then appears to be partly prepared by

(1) In Latin, *Auriculo-ventricularia*.

(2) In Latin, *Valvula interventricularia*.

transmission through the coats of the intestine, and again probably undergoes changes in its passage to, and its course through, the heart; and being diffused through the cavity of the body without being confined in blood vessels, it is probably distributed whither it is wanted by the muscular movements of the animal, and assimilated by what is termed imbibition.

ORGANS OF BREATHING.

IN the larger animals, the blood, by means of lungs, is exposed to fresh air, taken in by the mouth and nostrils, and through the windpipe passing into the lungs, where it gives off oxygen to the blood, and carries off carbonic matter from the blood¹. In insects there are no lungs, and consequently no air is taken in by the mouth (there are no nostrils) by breathing.

Fresh air, however, is as essential to insects as to other animals, though it enters their bodies in a different manner; since, instead of one, two, or three, openings for breathing, there are usually eighteen, generally very obvious in the larger caterpillars; their function being demonstrable by stopping them up with oil or grease, when the animal is soon suffocated and dies. The pores of plants seem to be similar².

These openings are termed spiracles³, and are of two sorts, simple and composite.



A spiracle magnified to show the lips open to inhale the air in breathing.

(1) See ALPHABET OF ZOOLOGY, page 85, &c.

(2) See ALPHABET OF BOTANY, 2nd edition, page 82.

(3) In Latin, *Spiracula*, improperly *Stigmata*.

The simple spiracles¹ are usually situated on the sides of the abdomen, a pair (one on the right and one on the left) to each ring, at the junction between the back arch and the belly arch. Like the nostrils in man and other animals, these spiracles are often beset with hairs, crossing and meeting, for the purpose of preventing the entrance of what might prove injurious, while the air may pass pure.


The position assigned them as well as their number, varies considerably, to suit the habits of the insect. In the maggots of some flies, for instance, which feed on greasy substances, or live in water, there is only one spiracle at the end of the last ring, which can be stretched out like a telescope in the air, while the body is enveloped.

The second and third rings of the body of caterpillars, answering to the mid and hind coxæ, having no spiracles in caterpillars, and the exact parts where they might be looked for being the place where the four wings are afterwards jointed, M. Blainville inferred that the wings, when expanded, in the moth and butterfly, were nothing more than spiracles. In this singular opinion, M. Audouin would have agreed, had it not occurred to him, that if the wings were to be considered spiracles, there would not be both wings and spiracles in the same insect at these four points, which upon examination he actually found, as in the cockroach and the mole cricket; and of course M. Blainville's fancy is thence annihilated.

The opening of these spiracles is surrounded by a ring, more or less circular, and somewhat elevated, which being contractile, may be considered as similar to the lips of the mouth.

(1) In Latin *Spiracula simplicia*.

The composite spiracles¹ are never placed in the abdomen, but exclusively in the fore corselet, and are (so far as is at present known) only two in number. These are very obvious in grasshoppers. They are composed of two horny pieces, which move outwards and inwards in the process of breathing, like a pair of folding doors, their movements being produced by two muscles.



The right side of the corselet in the cockchafer cut across lengthwise, to show the first layers of air pipes highly magnified. A, A fore corselet; B, B, middle and hind corselet; C, the first spiracle leading to D, the bulging of the air pipe; E, bulging of the air pipe from the third spiracle. 1, upper cephalic air pipe, a, its first branch; 2, lower cephalic air pipe; b, its first branch; 3, upper crural air pipe; 4, fourth trunk which goes also into the haunch; 9, the ninth trunk; 10 and 11, the tenth and eleventh trunks; 12, air pipes of the wing cases; 13, thirteenth trunk; e, small branches therefrom, forming cells; i, its point of junction² with the second trunk of the second spiracle; 14, fourteenth trunk of the first spiracle; f, g, h small branches going to the middle haunch; 15, middle crural air-pipes; k, l, n, the second trunk of the second spiracle; m, cells therefrom; o, third trunk of the same spiracle; p, p, cells therefrom; q, cross communicating branch; r, branch of the opposite side; s, reunion of two branches; t, large abdominal branch; u, a branch to the extensor muscle of the hind haunch; v, branch of the first arch; w, x, communicating branches; y, branches to the upper arches; z, branches to the organs of reproduction.

(1) In Latin *Tremæra* or *Spiracula composita*.

(2) In Latin, *Anastomosis*.

The spiracles convey the air which they inspire from without to a corresponding number of air pipes to be carried into the body, as the air inspired by man is carried into the windpipe and the lungs. The several air pipes which go from the spiracles end in a common pipe on each side, and these two common pipes may be termed the main air pipes¹. They run lengthways from the head towards the vent, and send off innumerable small branches to convey the air to the different parts of the body, somewhat like the branches of certain shrubs; the branchlets² interlace the membranes, penetrate the muscles, and extend through the legs and the wings.

Two sorts of these air organs have been distinguished, the one tubular or pipe-like, and another vesicular or cell-like.

The first sort, or air pipes,³ are composed of three distinct membranes, the outer and inner of which are thick, extensible, and of a cellular texture, while the middle one is formed of a gristly thread rolled spirally round in the manner of a corkscrew, and very similar to the spiral air pipes of plants. This gristly spiral thread is very elastic, in consequence of which the pipe is kept uniformly open, for even when it is compressed by the muscles, it immediately expands again. There is a similar mechanism of gristle in the human windpipe.


These air pipes have also been distinguished, from their situations, into arterial and pulmonary, the arterial being those which come directly from the spiracles; and the pulmonary, the two large pipes on each side of the body, (not always traceable), from which other arterial pipes branch off. The structure of both these is similar.

(1) In Latin *Tracheæ*.

(2) In Latin, *Ramuli*.

(3) In Latin, *Tracheæ tubulariæ*.

The second sort, or air-cells¹, are without the spiral gristle, being composed only of an outer and inner membrane. Consequently, those air cells, when not filled with air, must become flaccid by their sides collapsing. They are not in the form of pipes, but like cells or pouches, mutually communicating with each other through very short and simple canals. They never communicate directly with the spiracles, but



The breathing apparatus in the Praying Mantis, showing the numerous air tubes and air cells on each side.

(2) In Latin *Tracheæ vesiculariæ*.

receive their air from the air tubes. They are not found in all insects; but, when they occur, they appear to serve the purpose of reservoirs for air.

They vary also in number and in size. In the rose chafer, for example, they are very small and very numerous; but comparatively large in grasshoppers and crickets. In these insects, the air cells can be easily counted. It would be a difficult matter to inflate them with air, had there not been an ingenious contrivance to facilitate this, in a sort of ribs with which their sides are provided, discovered by Marcel de Serres, and consisting of small projections from the edge of each ring of the belly, and not distinct jointed members.

The air then is breathed by means of these various organs, and acts on the blood, or the fluid similar to blood, somewhat, it may be presumed, in the same way as it acts on the human blood, in the lungs,—oxygen being abstracted, and carbonaceous matter carried off¹.

The insects which live under water constantly or partially, have peculiar organs for decomposing the water or the air it contains in order to procure oxygen, which appears to be indispensable to life. Some of these water insects, indeed, come ever and anon to the surface, in the same manner as the water eft and the whale, to breathe the air. Others remain always under water.

ORGANS OF SENSATION.

It is well known that in man, the brain and spinal cord, with the nerves proceeding from them or con-

(1) This will be fully explained in the ALPHABET OF PHYSIOLOGY.

nected with them, are the organs of feeling or sensation, the brain being encased in the strong bones of the skull, and the spinal cord in the no less strong and peculiarly-jointed bones of the back. The nerves in insects differ much from those of man, and particularly in there being no peculiar structure like the skull, and the bones of the spine, to encase what may be termed the main stem, whether that be considered the origin or the receiver of the branches. This main stem in insects, however, which lies along the breast and belly, from the head to the vent, is protected by the inner breast-plate from being compressed by the gullet, as has been already noticed.

In man, the brain is distinguished into two parts, the spinal cord being a third; but besides these there is a system of numerous and extensive nerves, which is only connected with the three parts just mentioned by very small twigs, and is therefore considered as being partly independent of them¹. This partially independent system of nerves in man is termed the ganglionic system, and it is sometimes also called the great sympathetic nerve, or the intercostal nerve.

Now it is generally maintained by naturalists, that insects possess only this ganglionic system of nerves, and have no brain or spinal cord like that of the larger animals.

It may be well, before proceeding farther, to describe a ganglion, which is a knot or mass of nervous substance, at a point where two or more nerves meet, and appears to consist of two substances similar to those of the brain, while it differs from a nerve in being firmer in texture, redder in colour from a greater supply of blood, and covered with a membrane of

(1) See ALPHABET OF ZOOLOGY, p. 97.

closer tissue. The fibres or threads of the nerves which join such a nerve-knot or ganglion, become twisted within it, as Scarpa says, into a bundle, and threads from the several joined nerves unite to form a new nerve, which is always larger than any of those whence it has been formed. This constitutes a ganglion in man.


Instead of a brain, then, insects have generally in the head a double nerve-knot or ganglion, contained in a horny cavity larger than itself. This, for the sake of distinction, may be called the ganglionic brain¹; inasmuch as it differs from the ganglionic system in man by sending nerves to the eyes, the ears, and the mouth, while the former appears (so far as is yet understood) to supply nerves only to the heart, stomach, intestines, and other organs, whose motions are involuntary. The ganglionic brain also differs from the brain in man by being surrounded with powerful muscles which move different portions of it, whereas, there are no muscles to move the human brain. Besides the nerves to the eyes, the ears, the mouth, from the fore part of the ganglionic brain in insects, and two nervous films behind, difficult to detect, and probably running to the heart,—two thick nerves go off from the base, and after forming by their separation a sort of ring or collar, which, dipping down, embraces the gullet, they re-unite at what may be called the second nerve-knot or ganglion², below the gullet, whereas the brain is above the gullet.

In the same way, two nerves go off from the lower part of this second nerve-knot or ganglion, and re-unite at a third³.

(1) In Latin, *Cerebrum ganglionicum*.

(2) In Latin, *Ganglion secundum*.

(3) In Latin, *Ganglion tertium*.



Nervous system in the cockchafer on a horizontal plain magnified seven times. A, A, the two principal nerve-knots of the head; a, b, a, b, nerves of the ears; c, c, nerves of the eyes; t, swelling of the nerve as it enters the eye which is cut across to show its structure; d, collar of the gullet; c, e, f, two pairs of nerve-knots accessory to the first; g, root thereof; h, k, two nerve-knots of the vital organs; i, i, nerves from the upper of these; l, l, nerves of the third pair.

B, Nerve-knot under the gullet; m, m, nerves of the upper jaws;

a, n, their branches; *p, p*, nerves of the under jaws; *q, q*, their first branch; *r, r*, nerves of the third pair; *s, s*, back thereof.

C, Nerve-knot of the fore corselet. *a*, arms of this; *b, b*, first pair; *c, l*, nerves of the muscles; *m*, second pair; *n*, fork.

D, Nerve-knot of the mid corselet; *a*, nerves of the wing cases; *b, l*, nerves of the muscles; *m*, middle crural nerves.

E, Nerve-knot of the hind corselet; *a*, nerves of the wings; *b, r*, nerves of the muscles.

F, Nerve-knot of the abdomen; *a, e*, five pairs of nerves therefrom; *f*, terminal portion of the nerve-knot chain; *g, i*, three branches therefrom.

Two nerves go off from the third in a similar way, and thus, at intervals, a chain of nerve-knots or ganglia, united by double nervous cords, is formed along the belly of the insect to the vent.

The number of nerve-knots or ganglia varies in different species. Sometimes there is one for each ring of the body, and in other cases, not so many as this. The louse has only three, while the mole-cricket has nine, and the green field cricket has ten.

When one of these nerve-knots is carefully observed, it is found to be usually spherical or pear-shaped, sometimes flat, (in the gipsy-moth, the third is heart-shaped) consisting, like the ganglionic brain, of two lobes, not always of the same form, even in the same insect.

Besides the two main nerves, or double nervous chord which unites the several nerve-knots into a chain, there goes off from each, on both sides, small nervous branches, and these again divide into branchlets smaller and smaller, which are distributed to the gullet, the stomach, the air-pipes, and the muscles throughout the body.

Lyonnet counted forty-five pairs of these nerves, and two single ones in the goat caterpillar, making in all ninety-two, which is fourteen more than are found in man.

The double nervous chords uniting the ganglia are

tubular, and composed of two substances, the membrane forming the tube, and a sort of marrowy substance with which this is filled. It is probable, that the smaller branches are similar in structure.



The nervous system in a garden beetle (*Carabus*). *a*, the first nerve-knot or ganglion, with a large nerve on each side going off to the eyes; *b*, the second nerve-knot; *c*, the third nerve-knot; *d*, *e*, *f*, *g*, *h*, the other nerve-knots or ganglia.

It is not improbable that each nerve-knot may form the centre of feeling to the parts with which its nerves communicate; and if this be so, it will afford some explanation of the fact, which is no less singular than it is ascertained beyond question, that insects obviously

do not feel so much pain from wounds and injuries as larger animals. Hence it appears to be, that the abdomen of a wasp or a bee will continue to live, and thrust out the sting, long after it is severed from the body; and the head of a dragon-fly will eat as voraciously after it is cut off, as if it had to supply an insatiable stomach. The circumstances seem to disprove, in the most decided manner, the humane, though mistaken, opinion of the poet, so often quoted, that

"The poor beetle which we tread upon
In corporal sufferance feels a pang as great
As when a giant dies.

No giant could kick if his body was cut asunder, yet the bee stings in such circumstances; no giant could eat voraciously like the dragon-fly when his head is cut off, nor walk about without his head, as a common fly will readily do; nor after his bowels have been scooped out, as cockchafers often do.

May it not be that Providence has endowed insects with less acute feelings, in order to lessen their sufferings when preyed upon by birds and other animals, for whose food they appear to be mainly intended? I throw out this as a plausible conjecture.

Rationality and Instinct.

According to an ingenious and very plausible theory of Dr. Virey, which makes the brain and spinal cord the exclusive seat of intelligence and reason, insects must be devoid of these as they have no brain and spinal cord, at least, like those of quadrupeds. All their motives accordingly must be purely instinctive, that is operating involuntarily. "Among insects," Dr. Virey justly says, "there is no apprenticeship, no improvements, no variation in practice, no superadded invention." A

mason bee or a carpenter wasp is hatched from eggs laid by the mother who dies many months before her offspring acquires wings; yet this offspring will form the same ingenious nest as the mother had done without any other instruction than the impulse of instinct implanted by Providence. These are facts, of which hundreds of examples could be given. It would be well if Dr. Virey's theory or any other were as little liable to objection or refutation as these facts are¹.

THE SENSES OF INSECTS.

The difficulty of investigating the senses of insects is very great, owing to the want of analogy between the structure and circumstances of these little creatures and our own; a difficulty which is evidently much less in the case of quadrupeds.

The Sense of Touch.

The sense of *touch* has been by many supposed to reside in the organs I venture to call the ears, which have thence been termed feelers, but the evidence on which this rests is slight and unsatisfactory; for the bending the ears forward, and moving them in walking, seem to be for the purpose of listening.

The organs in question "being sheathed," says Professor Bonsdorf, "with a crusty shell, are therefore ill-fitted for feeling." "The conjecture," says M. Straus Dürkheim, "is founded upon facts imperfectly investigated, if not altogether false. I have made numerous researches on this subject, and I have never been able to satisfy myself that insects examine objects by feeling them with their antennæ. On the contrary, I have rarely observed these animals touch any thing with

(1) See ALPHABET OF ZOOLOGY, page 102.

these organs, and when this did happen, it appeared to be only by accident, and not at all from design. Many insects, besides, have their antennæ so short, that they would be obliged to stand erect upon their heads in order to come at the bodies which they might thus wish to explore, and for this their feet are certainly much better adapted."

I think that the numerous joints in the foot permitting it to bend round objects, and more particularly the soft-cushioned feet of some beetles and flies, and also the feelers on the under jaws, and the under lip, may be more plausibly considered the organs of touch; while the hairs, and long bristles, and spines of some caterpillars, are evidently connected with the sense of touch.

The Sense of Taste.

The younger Huber describes the tongue of the ant as a minute, conical, fleshy, yellowish organ, which is alternately pushed out and drawn in. M. Straus Dürckheim says the tongue of the cockchafer is small, fleshy, and moveable; but he is more inclined to consider as the organs of taste, what he terms the jaw brush, consisting in beetles of a tuft of thickly set stiff hairs around the grinding surface of the upper jaw. This I consider as pure conjecture, as to ascribe taste to the feelers.

Wherever the organ be situated there can be no doubt that insects possess the sense of taste, were there no other facts to prove it than that the gnat, the midge, the bug, and the flea will severely attack some individuals, particularly women and children, and not molest others.

The Sense of Smell.

The Swedish naturalist, M. Clerck, in 1764, confi-

dently asserted the organs which I have called the ears to be those of smell, because the rose chafers open their plates on alighting on flowers, as horses expand their nostrils to the cool air after a hard gallop. Bergmann with the same view referred to the cuckoo fly *smelling* as he thought with its long ears into the nests of wasps. It is singular that Réaumur and Lyonnet had a similar opinion, which as Straus Dürckheim justly remarks, "is supported by no fact either anatomical or physiological."

Most insects possess an exquisite sense of smell, but as they do not, like quadrupeds, breathe through nostrils and do not possess them, we are naturally led with Baster to suppose the organ of smell to be in the spiracles:—M. Huber thinks in a small spiracle under the tongue, while M. Dumeril appears to think that the whole lining of the air pipes throughout the body is the organ of smell—evidently a gratuitous assumption highly improbable; for as M. Straus Dürckheim remarks, the sensation must depend on the expansion of the nerves of smell.

M. Marcel de Serres is strongly of opinion that the organ of smell is in the interior of the feelers, which are lined with a moist membrane furnished with nerves and communicating, at least in crickets and grasshoppers, with the external air by a small opening. He can see no good reason, nor can I, for supposing what I have termed the ears to be organs of smell; and he thinks the absence of nerves in the spiracles, opposed to the opinion of Baster, above stated.


The Sense of Hearing.

For the brief reasons assigned under touch and smell, and for others deduced from dissection and experiment, I have ventured to call the *Ears*, two horn-like organs, always situated near the eyes, to which

various incongruous functions have been assigned. As I have little doubt these organs will one day be proved to be ears, I think it will direct attention more decidedly to them by at once terming them ears, than by leaving them open to all sorts of crude fancies, so easy to form, but so detrimental to correct inquiry. Before stating in brief my reasons for this opinion, it may be well to show the structure of the ears.



Structure of the ears in the cockchafer placed before the eyes and magnified eight times. *a*, *b*, the three first joints of the ears cylindrical; *c*, the seven joints at the tip short and in form of leaves; *h*, a pertractor muscle; *i*, an elevator muscle; *k*, *k*, *k*, abductor muscles; *l*, *l*, *l*, adductor muscles; *o*, large nerves arising from the first nerve-knot or ganglionic brain *λ*, within the head.



Air pipes in the ears of the cockchafer magnified eight times. *a, a*, the main branches from which spring the ear branches, *b, b*.

Professor Bousdorf from some rather loose and vague experiments, such as making loud sounds and observing insects in consequence move the organs in question, concluded them to be ears; an opinion also held by Göze and Christ.

When I was writing "INSECT MISCELLANIES"¹ at Havre de Grace, in 1829, I placed a green grasshopper remarkable for its long ears upon a table, and on rustling a bit of paper under the table out of view, it immediately bent one or both of its ears in the direction of the sound, just as a rabbit would do if similarly alarmed. Of many sounds, such as the beating of a watch and the ringing of glasses, it took no notice. I repeated these experiments in every possible way on many species with similar results. M. Straus-Dürckheim published in the same year very similar remarks, though he does not particularise any experiments similar to mine. He thinks the numerous joints in the ears intended both to prevent their being easily broken by accident, and "to perform the same functions as the chain of small bones in the human ear, inasmuch as they form a similar chain and transmit the vibrations of the air to the auditory pulp."

It is not a little singular that M. Bonnet denied

(1) See Chapter IV.

insects to have any sense of hearing—an opinion also maintained by Linnæus.

The Eyes and the Sense of Vision.

In the larger animals, there is only one sort of eyes, but insects have two sorts, very different in structure, which have been called simple and compound, there being always two compound eyes¹, placed near the base of the ears on each side of the forehead or face; but the simple eyes² vary in number, and are placed higher up, usually on the crown of the head, as in the bee, where the simple eyes are three in number, and placed in a triangle.



The upper portion of a bee, showing the two compound eyes with their facettes at the base of the jointed ears; together with the three simple eyes in the form of a triangle on the crown.

We do not meet with simple eyes in all insects, but no adult insect is without compound eyes. The number of simple eyes is usually three. Each is composed of, 1. an outer membrane, hard, transparent, and formed of a single piece³; 2. a layer of a viscus or clammy substance⁴, immediately behind the first, which determines the colour of the eye, being black in bees, white in crickets and grasshoppers, and red, yellow, or green in some caterpillars; 3. a rather thick membrane⁵,

(1) In Latin, *Oculi compositi*.

(2) In Latin, *Ocelli*, or, very objectionably, *Stemmata*.

(3) In Latin, *Cornea*.

(4) In Latin, *Pigmentum*

(5) In Latin, *Choroides*.

seemingly composed of a tissue of net-work, whose meshes are very closely set; and 4. a nerve¹ from the first nerve knot or ganglion of the head, very small, which penetrates to the inside of the outer membrane, where it spreads out.

Compound, composite, shagreened, or faceted eyes, are so named from being made up of a great number of minute eyelets, not unlike the six-sided facettes of crystals. In the cockchafer, these, according to Straus-Dürckheim, are 8820 in number. They are always immoveable, differing in this from the eyes of larger animals. Each of the little eyelets or facettes is very similar in structure to one of the simple eyes already described. According to Professor Müller, of Bonn, the composite eye of the dragon fly may be divided into two parts; one above and behind of a reddish colour, with the eyelets twice as broad as those of the other in front and below, which is greyish.

When the whole composite eye is cut into, we find, 1. the outer transparent membrane (*a*); 2. the larger or clammy-coloured matter (*b*); 3. a broad belt, orange-coloured before, and black in front (*c*); 4. a second belt within the first, deep black (*d*); 5. the ganglion of the nerve, which, when slightly pressed, is seen to be composed of rays of fibres, or threadlets (*e*), one of which probably passes to each eyelet.




See also the figure in page 83.

(1) Technically, *Optic nerve*.

The cellular tissue, and the clammy-coloured substance, are found wanting in some species of night insects; and, according to Treviranus, there is in the cockroach, (a night insect,) behind the outer membrane, a mass of a dark violet colour, composed of numerous little pyramids, upon which the nerve is spread in the form of fibres. The eyes of many night insects shine in the dark.

The vision of insects is much better understood than that of other senses. As I have already described the simple and compound eyes of insects, so far as their outward structure is concerned, I shall now call attention to the nerves of the eyes. ¹

In insects which have large eyes, these nerves are exceedingly large, bulging out after they go off from the ganglionic brain into considerable knobs. In the stag-beetle, in which these are pear-shaped, so many minute branchlets go off to the eye, (one probably to each facette,) that it is not possible to count them. In the hive bee the knobs of these nerves are kidney-shaped, and so much larger than the brain itself, that they might lead an indifferent observer to suppose they were actually the brain.



The head nerves in the bee. *a*, the first nerve knot or ganglion, with its forked division below; *b*, the small nerves of the head; *c, c*, the two large nerves of the eyes.

(1) In Latin, *Nervi optici*.

It is remarkable that the eyes of insects are supplied with large air pipes, arising from the main air pipe in the head; one rather large, surrounding the eye, and many others going off from this and dividing into equal angled triangles, as has been minutely described by Marcel de Serres.

I have proved by experiment that M. Latreille was mistaken in supposing a small black ant, common in France, to be blind: I found it on the contrary very impatient of light.

ORGANS OF REPRODUCTION.

UNLIKE snails and worms, among which there is no distinction between the males and females, insects have the two sexes as distinct as the larger animals, and in many respects are similar to birds, so far as pairing is concerned,—that is, a single male associates with a single female. Exceptions to this general rule occur among hive bees, and the white ants of warm countries; a single female, called the queen, being attended by many males, while the young are nursed by a peculiar class, (not exactly females, but more like these than like males,) called nursers or workers. The reverse takes place among poultry and black cattle, one male being attended by many females. Among ants, wasps, and humble bees, several individual females in the same hive are attended by several males.

Insects differ from birds and many other animals in the male taking no share whatever in providing for the young, either before or after pairing.

The male is generally smaller than the female, sometimes, but not always, brighter or differently coloured; is distinguished in moths and gnats by the ears being

feathered or ornamented, while those of the female are plain; and in grasshoppers and cuckoo-flies by wanting the ovipositor. In some insects the male has ample wings, when the female has none, or very minute ones, not adapted for flying.



The Vapourer moth, the female with plain ears and with very short wings; the male with feathered ears and large wings.

The male insects are, in most cases, more restless and wandering than the females, and of course more frequently seen, differing in this from spiders, the males of which are seldom or rarely seen.

All female insects have eggs, which, in a few cases, are hatched within the body, but are generally laid in such places as the young may readily find food when hatched; the mother, in most instances, dying soon after the laying, and of course the young have from the first to shift for themselves.

These are ascertained facts which cannot be questioned, though they do not accord with the popular error of insects being generated by putrefaction, and by blighting fogs or winds, much less the so called philosophical theory of their being generated by some sort of mysterious chemistry.

This theory was supposed to be unanswerably supported by the multiplication of microscopic animalcules in water, which could not, on account of their

minuteness, be traced to their parents. This, however, though apparently impossible, has recently been done by Professor Ehrenberg of Berlin, who, by putting the animalcules in coloured fluids, succeeded not only in discovering their eggs, and the hatching of these, but in tracing them before they were laid in the egg-organ of the mother¹.

M. Ehrenberg was no doubt assisted in this by the circumstance, which generally holds good, of the eggs of insects and small animals being proportionably much larger, compared with the mother, than in birds.


From this circumstance, the eggs of insects may, in many cases, with a little care, be traced in the egg-organ of the mother, as has been done by M. Straus-Dürckheim in the cockchafer, and by Swammerdam in a much smaller insect—the louse.



Egg organ in the louse magnified; *a*, the right branch converging; *b*, the left branch diverging; *c*, an egg in the egg tube ready to be excluded; *d, d*, the fountain which furnishes the glue.

When an egg is laid by an insect, it is, except in a very few cases, glued to the place where it is laid by a sticky fluid provided for that purpose, and discharged along with the egg, or immediately after it; the vessel producing which fluid in the louse may be seen in the

(1) In Latin, *Ovar. u. u.*



Male reproductive organs in the blister beetle.

figure. The louse glues its eggs to hairs; moths and butterflies glue theirs on the bark or leaves of plants and trees, where they usually remain through the winter, to be hatched the following spring.

It is thence obvious that the eggs of insects, thus firmly glued, cannot, as has been fancied by the unlearned, as well as by philosophers, float about in the air, than which, indeed, they are always much heavier.

Several species of insects, instead of gluing their eggs in this manner, place them in nests, as is done by the bees and wasps, appropriately called miners, masons, carpenters, or upholsterers, according to the processes they pursue in building their nests.

Other species dig holes in the ground, or cut out grooves in wood, in which to lay their eggs, and are provided with curiously-constructed instruments for that purpose, such as the grasshopper, whose digging instrument is like a scymitar; the saw flies, whose instrument is like a double saw; and the gall-flies, whose instrument is like an awl.

Several species, well termed cuckoo-flies, thrust their eggs into the nests or the bodies of other insects, or animals, and have appropriate boring instruments for this purpose.



Instruments for depositing eggs; *a*, that of a cuckoo fly; *b*, that of a grasshopper; *c*, that of a saw fly.

This is indeed one of the most singular circumstances in the economy of insects, and frequently leads to no little disappointment, when collectors feed caterpillars for the purpose of breeding moths from them; for, instead of the expected moth, a brood of cuckoo-flies, whose maggots had been feasting on the body of the caterpillar, make their appearance, and immediately pair and set out upon excursions for the discovery of other caterpillars, into which they may thrust their eggs.

GROWTH OF INSECTS.

I HAVE here selected the word "growth" in preference to *development*, both because it is shorter, more English-like, and less startling to beginners, and because *development* has recently become one of the catch words of an absurd theory, and has been twisted from its true meaning, by applying it comparatively to two or more animals, instead of confining it, as it always should be, to an individual. *Metamorphosis*, or *Transmutations* is still more objectionable, where similarly applied, and I have elsewhere used "Transformations," as somewhat less so.

EGGS OF INSECTS.

INSECTS' eggs are not all of an oval form like those of birds, but some are like a pear, some like an orange, some like a pyramid, and some like a flask.



Various shaped eggs of insects magnified.

The eggs of the gnat, for instance, may be compared, in shape, to that of a powder flask, and the mother gnat lays about three hundred at a time. Now each egg, by itself, would sink to the bottom of the water; yet the gnat puts the whole three hundred together in the form of a little boat, and in such a way, that they will all swim on the surface of the water; and a very curious way she has of managing this.

Like other insects, the gnat has six legs. Four of these (the four fore-legs) she fastens to a floating leaf, or to the side of a bucket, if she is on the water contained in one. Her body is thus held level with the water, except the last ring of her abdomen, which is a little raised. This being done, she begins to make use of her other two legs, (or hind legs) and crosses them in the shape of the letter X. The open part of this X, next to her tail, serves as a kind of scaffolding, to support the eggs she lays, until the boat is formed. Each egg, when laid, is covered with a kind of glue; and the gnat holds the first laid egg in the angle of the X until the second egg is laid by its side, and glued to it; she then glues another egg to its other side. All these stick together thus * * *, making a kind of triangle, or figure of three, and this is the beginning of the boat. Thus she goes on, piling egg upon egg, always keeping the boat in proper shape by her useful hind-legs. As the boat grows in size she pushes it from her by degrees, still adding to the unfinished end next to her body. When the boat is half built, her hind-legs are stretched out thus =, the X or cross form is no longer wanted, and she holds up the boat as cleverly as if it was done with two out-stretched arms.

The boat is at length completed, and an excellent boat it is, quite water tight. For though it is very small and delicate, yet no tossing of the waves will sink it; and nothing can fill it with water, or turn it upside down. In fact, the glue with which it is covered prevents it from ever being wet. Even if the boat be pushed down to the bottom of the water, up it comes again quite dry: so that it is better than the best life-boat that has ever yet been invented¹.

(1) Working Man's Companion, Cottage Evenings, page 54.



The eggs of insects are not, like those of birds, always smooth; but are sometimes ribbed, and sometimes tiled, or otherwise sculptured or carved on the outside.

The shell of an insect's egg, is rarely or ever brittle, like that of a bird, but composed of a tough membrane which, in some instances, can be stretched out, as appears from the eggs of ants and some other insects growing considerably larger in the process of hatching.

The mother insects, usually dying before their eggs are hatched, do not sit upon them like birds, except in the singular instance of the earwig, which, from the proceedings of one kept by me in a glass, in March, 1832, appears to attend more to shifting the eggs about to places where they may receive moisture, than any thing like hatching by covering them. Ants shift their eggs according to the changes of the day and night, and also of the weather, placing them near the surface of their nests when it is warm and dry, and deep down when it is cold and wet.

In consequence of being exposed to the same temperature, all the eggs of any particular species, in any given district, are hatched exactly at the same time, or at most within a few days; and when such eggs are numerous, an immense number of caterpillars make their appearance all at once on plants and bushes, and give rise to the notion that they are brought by winds, or generated by what is called *blighting* weather; though this is as absurd as to say the wind could bring a flock of cattle, or that the *blight* could generate a flight of sparrows or rooks, without eggs to hatch them from.

By looking carefully on the bark of currant and other bushes, or on the back ribs of gooseberry leaves, the eggs may be found sometimes in patches, sometimes

in rows, whence the caterpillars are hatched that creep into the buds, or stream over the leaves and devour them.

INFANCY OF INSECTS.

THERE is no single English word which will apply to every insect just after it is hatched, and while it remains in what may be called its state of infancy ; but there are several English words which apply to this state in different sorts of insects, as I shall now explain.



Insects in the infant state ; *a*, caterpillar of a saw fly ; *b*, grub of a ladybird ; *c*, maggot of a cheese fly magnified.

The word "CATERPILLAR"¹ is applied to the creatures which are, with a few exceptions, shaped like an earth-worm, and of various colours, but most commonly green ; sometimes smooth, sometimes studded with short or with long hair, and sometimes with a sort of thorns, and having exactly six legs always placed on the corselet and furnished with claws, while they have from two to sixteen clinging feet² without claws, always situated on the rings of the abdomen. All caterpillars are hatched either from the eggs of

(1) In Latin, *Eruca*.

(2) In Latin, *Prehensores*, objectionably *Propedes*.

butterflies, moths, or saw flies, with four wings, and, when full grown, they become butterflies, moths, or saw flies, like their parents.

Among the caterpillars most known and most destructive, are gooseberry and willow caterpillars, which become saw flies, as does the small caterpillar, erroneously called the turnip fly¹. Other well known caterpillars are the large caterpillars which feed on cabbage, and become white butterflies, and small gregarious caterpillars which destroy hedges and fruit trees, afterwards becoming moths, as do those which roll up the leaves which they eat, or creep into buds, constituting the well known "worm i' the bud." Other caterpillars, which become small moths, devour grain in granaries, the wood of currant and willow, or other trees, garments of woollen or silk and furs, and insects and other animals kept in cabinets.

The word "GRUB" is not quite so definite as caterpillar, and is often applied popularly, but erroneously, to both caterpillars and maggots. In precise language, a grub is a creature hatched from the egg of some sort of beetle or weevil (which is a beetle with a longish snout). A grub has always exactly six feet on the corselet, and never any clinging feet on the abdomen like caterpillars. The body of a grub is also, with a few exceptions, more clumsy than that of a caterpillar, the general colour being white, yellow, or brown, never, I believe, green; and the head usually some darker colour, as blackish or dark brown.

The best known grub is that of the cockchafer, which takes three years to arrive at its full growth, and devours the roots of grass; those of the corn weevil, which do great injury in granaries; and those

(1) There is a small jumping beetle also called the turnip fly.

of the weevils that feed on nuts, and on the buds of apple and other fruit trees, as well as on the roots of cabbages and turnips, producing knobs on them. The meal-worm and the wire-worm are also grubs, as well as those which worm-hole furniture, gnaw bacon, and destroy the bark of trees. All these become beetles in their adult state, which in turn produce a fresh laying of eggs from which other grubs are hatched.

The term "MAGGOT" is more precise than grub, though some maggots are erroneously called grubs. Maggots are never produced from the eggs of butterflies, moths, saw flies, beetles, or weevils; but always from the eggs of two-winged flies, or from bees or wasps. A maggot differs from a caterpillar or a grub in having no feet, and from an earthworm in never being of a dull red or dingy green colour, but usually white, greyish, often transparent, so as to show the intestines, and, in the case of the water maggots, called blood-worms, of a bright blood red.

The best known maggots are those of the blow-fly, which live on meat, either fresh or putrid; those of the bot fly, producing the disorder called bots in horses; those of the cheese-fly, called jumpers, or hoppers, and sometimes erroneously mites, (mites not being insects at all, but ranking with spiders;) and those of the crane flies, which destroy grass and corn fields, often improperly called the grub. The wheat fly, as it is called, and the Hessian fly, which have proved so destructive, are both grubs of small flies, not unlike gnats.

Maggots are popularly called mawks in the North, and gentles in the South, and very often worms; as in the instance of the blood-worm, which is the infant

state of a small gnat fly¹, and which has been shown, by Mr. J. Fennel, to construct a covering for itself of sand and slime.



Blood-worm forming its covering.

These three sorts—caterpillars, grubs, and maggots—are, by modern naturalists, called by a Latin word, which means a mask or a phantom² because Linnæus took a fancy to suppose them only insects in a mask, which when they had thrown off, they were, of course, unmasked. I think the term objectionable and improper, though it is often convenient for want of a better.

These infant insects, indeed, often throw off their skins, or moult, as it may be termed, during their growth, the old skin partly splitting and sloughing off; but this old skin is no more a mask than the feathers of a bird, which are moulted once or twice a year, can be called a mask.

Like most young animals, all caterpillars, grubs, and maggots, eat voraciously, as it is necessary to supply nutriment for their increasing growth.

(1) Figured in ALPHABET OF ANGLING, page 38.

(2) This Latin word is *Larva*.

ADOLESCENCE OF INSECTS.



Adolescent insects; *a*, the peacock butterfly; *b*, the blue-bottle fly; *c*, the cockchafer; *d*, the dragon-fly.

THE English word adolescence, though derived, like infancy, from the Latin, is well understood as applied to the period of life between infancy and full growth. Except this, there is no term which occurs to me as applicable to the stage of insects succeeding the one just described. I have only adopted it here, however, not because I think it good, but because I cannot find one more suitable.

In this stage, insects differ more remarkably from the larger animals than almost in any other particular; for, while the latter remain active and continue, as before, to eat and grow, the former in few instances do either, but remain in a very singular state of torpor.

Preparatory to going into this state, many insects make a sort of nest, called cocoon¹, to lie in; others hang themselves up by the tail, by a very ingenious process; and some remain active and move about, as in the case of bugs, locusts, grasshoppers, crickets, and dragon-flies.

Besides the insects which make a cocoon, many of those which do not, upon moulting their last skin in

(1) In Latin, *Incunabulum*.

the caterpillar, grub, or maggot state, acquire a covering, in some cases horny and tough, in others thinner and more transparent. In this envelope they remain, till the soft juices accumulated in their previous state become duly assimilated to their proper organs or members, and form the horny skin of the body, together with the legs and wings.

In butterflies, such as the peacock and the alderman, this covering appears shining, as if gilt, and a Latin¹, and also a Greek² word, implying gilding, was given to butterflies in this state; and thence these words were extended to instances in which there was no gilding, and the terms, from frequent use, may now be considered half naturalized as English words, particularly the term *chrysalis*.

Those with gilt coverings, as is the case with most butterflies, are in this state of a longish form, with several projecting corners and angles, with no appearance of feet or wings. Moths, again, in this state, are usually longish and tapering, with distinct rings and spiracles for breathing along the sides. Two-winged flies are in form of an egg, except being equal in thickness at each end. Most beetles and gall-flies have no case-covering, but remain with their legs folded closely over their breast. It has been already mentioned that bugs, crickets, grasshoppers, and dragon flies are not thus laid up in case-coverings, nor do they rest with their legs folded, but walk actively about.

Insects, when laid up in these case coverings, were fancied by Linnæus to resemble infants bewrapt in swaddling bands, once common all over Europe, and a Latin word³, indicating this, was thence applied;

(1) The Latin word is *Aurelia*.

(2) The Greek word is *Chrysalis*.

(3) The Latin word is *Pupa*, better *Pupum*.

and to those not laid up, but remaining active, the fanciful term, *nymph*. The latter is objectionable, because it can only be applied to female insects, while it has been indiscriminately used for both male and female. The former, though almost naturalized in English, is liable to the same objection, and ought, therefore, to be in the neuter gender. So far as this term means an infant, it is correct, but the Romans used it both for a baby, a doll, and a girl before arriving at womanhood; and in this latter sense it may be used, if no better term occur.

ADULT STAGE OF LIFE IN INSECTS.

IN the same vein with the fanciful resemblance of a caterpillar to a masked insect, and of an adolescent insect to an infant wrapt in swaddling bands, Linnæus fancied that the adult insect, now unmasked and unswaddled, resembled a picture, image, effigy, copy; pattern, or representation of an insect; and thence adopted a Latin word¹ implying this, which is still used by modern writers, though evidently both far-fetched and incorrect, as it plainly means not the insect itself, but a statue or spectre of it. The disciples of Linnæus, however, accustomed to the art of inventing fanciful meanings for words, and of mystifying the plainest facts, will, no doubt, maintain that *Imago* does not mean an image, but the insect itself; in the same way as those of the modern schools will maintain that what they fancifully call *types* are not like any other known "types," the moulds, models, or dies struck for the purpose of forming figures, nor are they figurative representations, but the very animals

(1) The Latin word is *Imago*, whence our English word *Image*.

themselves. It is thus that fancy and inaccuracy take the place of facts, and do incalculable injury to genuine knowledge.

I would have had less objection to adopt, as I have elsewhere done, the term *perfect*, as applied to adult insects, had it not been in recent times grossly abused in theoretical comparisons of one animal with another, —a beetle, for instance, being termed *perfect*, because several pieces of its mouth are moveable; and a gnat or a butterfly *imperfect*, because the pieces of the mouth are joined into the form of a sucking pipe. The distinction of the two forms is certain and undeniable; the terms employed are no less impious than insulting to common sense.

After remaining in the adolescent stage for a due length of time, in some instances only a week or two, —very commonly for several months, particularly the winter months,—the case covering, when there is one, or, when there is none, the outer skin, is thrown off, and the insect emerges in its adult or full grown state.



The Twenty-plumed Moth magnified.

For a few hours after its emergence, it is usually somewhat moist, the wings being unexpanded, and, as it were, crumpled up, and the body not so distinct in

its outline as it afterwards becomes. In some beetles also, such as ladybirds, the colours are pale, and the first pair of wings, which form the wing cases, are soft, and of a pale uniform colour, without the spots afterwards seen on them.

It is now that the air tubes distributed through the body, and particularly through the wings, perform the important office of inflation; and if a main wing-rib be then cut across near the root, the part of the wing to which it goes ever after remains shrivelled.

Those who are little acquainted with insects, are exceedingly apt to think they grow like other animals, and from this cause commit many mistakes, not perhaps of great moment, but which it may be proper to rectify by examples. The most common British butterflies are those which are white, and all are usually looked upon as of the same species, differing in nothing, except, perhaps, in the size, the latter being erroneously ascribed to difference of age. But the fact is, that there are a considerable number of species of our white butterflies, and probably more varieties even of these than have yet been ascertained or described. It is certain, indeed, that butterflies do not, like the larger animals, increase in size as they grow older; for every individual, from the moment it becomes a butterfly, continues invariably of the same size until its death. Butterflies, indeed, seldom live longer than a few days, or, at most, a few weeks, and during this time they eat nothing except a sip of honey: and since this is so, it would be absurd to expect that they could increase in size. It must not, however, be understood from this, that the same species will always measure or weigh precisely the same; for though this will hold as a general rule, there are many exceptions, arising from the accidents the caterpillar may have suffered, from

which an individual butterfly originated. It is only during the caterpillar state that the insect eats voraciously, and grows in proportion; and if it is, during this stage of existence, thrown upon short allowance, it cannot acquire the standard magnitude, and the butterfly will be dwarfed from the first.

The same remarks with respect to growth, apply to insects of every kind; and the fact cannot be better exemplified than in the uniformity of the house-fly, among which scarcely an individual in a thousand, will be found to differ a hair's breadth in dimensions from its fellows. The smaller flies that sometimes mingle with the common house-fly, are those that come from the maggots in cheese.

Another very common instance of such mistakes occurs in gnats and mosquitoes, there being a very great number of species, all popularly designated by one or other of these names. Two of the mosquitoes of Brazil are here figured, to shew the great difference in form and structure.

Fig. 1.



Fig. 2.



Mosquitoes of Brazil. Fig. 1. *Culex molestus*.—*a*, natural size;—*b*, magnified. Fig. 2. *Simulium pertinax*.—*c*, natural size;—*d*, magnified.

Few insects, after arriving at the adult state, live more than a few days, or at most a few weeks: some live for a few months, but this is an exception to the general rule, of which we have examples in the female wasps and female humble bees, which after pairing very late in autumn, live through the winter, after all the males and workers have perished, and found new colonies the ensuing spring. It is not, so far as I know, ascertained beyond dispute, that the primrose butterfly and others, which are seen in early spring, have lived through the winter in the adult or in the adolescent state, though the latter is most probable. I have in vain made repeated trials to keep the adult insects, which uniformly died in a few weeks.

SYSTEMATIC ARRANGEMENT OF INSECTS.

THE only use of a systematic arrangement of insects or other natural productions, appears to me to be its convenience of bringing things together in some logical order, both to aid the memory in remembering, and the judgment in comparing and deciding upon, agreements and differences. A system, in this point of view, is similar to the frame work of a cabinet, into the partitions of which many little facts may be stored and dove-tailed, that would otherwise be scattered through the memory at random at the great hazard of being lost.

Such systems, however, since the time of Linnæus, who set the baneful example, have been considered the sole end and aim of study, and are even preposterously represented as a high branch of philosophy, though, viewed in this light, they appear to me no higher nor more important than the play pebbles, which a child may be seen to amuse itself in classing in rows or circles, according to size, form, or colour.

Nay,—viewed in the light of a philosophical study, they are worse than trifling,—they are decidedly injurious, by leading to serious errors, such as that of sucking insects, having “jaws totally useless, can do no injury to the agriculturist,” which has been asserted and published within the last two years, by a fanciful theorist, with the ravages of the hop-fly and the bean dolphin staring him with flat contradiction in the face¹.

(1) The same theorist has since proposed to prove the doctrine of the TRINITY, by what he calls a Trinarian System of Animals.

In Botany again, what is preposterously lauded as the *Natural* system, is asserted to lead at once to a knowledge of the qualities of a plant by merely ascertaining what is called its *natural* order; in the face of the glaring facts of the wholesome potato with the poisonous deadly night-shade, and tobacco, being found in one of these so-called *natural* orders; and the wholesome bread-fruit tree, the fig, and the mulberry, with the most poisonous known vegetable, the upas, in another of these *natural* orders¹.

Confining ourselves to insects, all sorts of blunders are committed by those who undertake to display what they term the natural system. Moths, for example, are said to be distinguished from butterflies by flying only during the night, though there are numerous examples of day-flying moths; and even groups are actually termed diurnal by the very theorist, who set out by telling us they are nocturnal. A nocturnal-diurnal moth can only, I should think, be imagined to be of pure Hibernian origin.

There is no system, indeed, without some such striking error either of omission or commission, and hence we must be contented to make the best we can of their imperfections.

I am most particularly disposed to object to the very fanciful and theoretical terms *Kingdom, Tribe, Family*, and the like, not only as being founded on no obvious ground, there being no proper king, chief, or head, in the divisions so termed, but actually involving serious hypothetical errors, contradicted by well-known facts. The terms "Class," "Order," "Rank," "Species," "Variety," from having some basis to go upon, are

(1) This is more fully explained in the "ALPHABET OF BOTANY," 2nd edition, page 181.

much less objectionable; though it ought always to be borne in mind, that such arrangements are only exercises in logic, and depend far less on the things arranged, than on the views of the arranger. The misconception of this fact has led to innumerable idle and ignorant discussions about natural and artificial systems.

Rejecting then the more recent and partially fashionable systems as hypothetical, that of Fabricius as complex and unnecessarily difficult, and that of Linnæus as wanting the grand requisite of distinctness, from bringing too many species under a genus, I think I cannot propose a better than that of the distinguished Swedish naturalist, Baron De Geer, from whose magnificent work on insects, perused with avidity in my younger years, I derived that taste for the study which increases with every coming year. In giving an outline of this system, I shall take the liberty of making several alterations, which I think will render it more easy in assisting beginners to arrange the insects which they may collect, or may wish to study.

In the ALPHABET OF ZOOLOGY where a general arrangement of animals is given, it may be seen that insects stand first in the third class of jointed animals¹ with spiders, crustaceous animals, and ringed worms following them in the same class.

(1) In Latin, *Articulata*.

CLASSIFICATION OF INSECTS.

INSECTS have exactly six feet, and the body divided more or less distinctly into head, corselet, and abdomen. Spiders, mites, scorpions, crabs, lobsters, shrimps, wood-lice, centipedes, leeches, worms, are, therefore, not insects. Insects may be

DIVIDED INTO

- I. *Winged Insects*¹, with four or two wings.
- II. *Wingless Insects*², with no apparent wings.

WINGED INSECTS¹.

DIVIDED INTO

1. *Four-winged Insects*³, with all the wings membranous.
2. *Wing-cased Insects*⁴, with two membranous wings under two wing-cases, more or less thick or hard.
3. *Two-winged Insects*⁵, some with poisers at the base of the wings, and some without.

FOUR-WINGED INSECTS³.

DIVIDED INTO

- I. SCALY-WINGED INSECTS⁶, with the mouth composed of a spiral sucker, and comprehending,
 1. Butterflies⁷—

a. Papilionians ⁸ .	c. Lycænians ¹⁰ .
b. Nymphalians ⁹ .	d. Hesperians ¹¹ .

(1) In Latin, PTEROTA.

(2) In Latin, APTERA.

(3) In Latin, TETRAPTERA.

(4) In Latin, ELYTROPTERA.

(5) In Latin, ANELYTRA.

(6) In Latin, LEPIDOPTERA.

(7) In Latin, *Papiliones*.

(8) In Latin, *Papilionidæ*.

(9) In Latin, *Nymphalidæ*.

(10) In Latin, *Lycænidæ*.

(11) In Latin, *Hesperidæ*.

2. Moths¹, divided into

- | | |
|--|--|
| <i>a.</i> Hawk Moths ² . | <i>d.</i> Geometrians ⁵ . |
| <i>b.</i> Afternoon Moths ³ . | <i>e.</i> Evening Moths ⁶ . |
| <i>c.</i> Owl Moths ⁴ . | |

II. HAIRY-WINGED INSECTS⁷, with the wing ribs more or less hairy, and comprehending,

- | | |
|--------------------------------|---------------------------------|
| 1. Salmon Flies ⁸ . | 3. Caddis Flies ¹⁰ . |
| 2. Stone Flies ⁹ . | |

III. LACE-WINGED INSECTS¹¹, with the wing ribs netted like lace, and comprehending,

- | | |
|---|-----------------------------------|
| 1. Pearl Flies ¹² . | 4. Dragon Flies ¹⁵ . |
| 2. Timber Lice ¹³ , mostly
without wings. | 5. Day Flies ¹⁶ . |
| 3. Ant-lion Flies ¹⁴ . | 6. Scorpion Flies ¹⁷ . |

IV. YOKE-WINGED INSECTS¹⁸, with the first and second pair of wings unequal in breadth, and yoked together, comprehending,

- | | |
|-------------------------------|---------------------------------|
| 1. Gall Flies ¹⁹ . | 5. Wasps ²³ . |
| 2. Ruby Tails ²⁰ . | 6. Cuckoo Flies ²⁴ . |
| 3. Ants ²¹ . | 7. Saw Flies ²⁵ . |
| 4. Bees ²² . | |

(1) In Latin, *Phalæna*.(3) In Latin, *Bombycina*.(5) In Latin, *Geometrina*.

(7) In Latin, TRICHOPTERA.

(9) In Latin, *Leptocerida*.

(11) In Latin, NEUROPTERA.

(13) In Latin, *Psocida*.(15) In Latin, *Libellulida*.(17) In Latin, *Panorpida*.(19) In Latin, *Cynipida*.(21) In Latin, *Formicida*.(23) In Latin, *Vespida*.(25) In Latin, *Tenthredinida*.(2) In Latin, *Sphingina*.(4) In Latin, *Noctuina*.(6) In Latin, *Tortricina*.(8) In Latin, *Phryganida*.(10) In Latin, *Tinodida*.(12) In Latin, *Periida*.(14) In Latin, *Myrmeleonida*.(16) In Latin, *Ephemerida*.

(18) In Latin, HYMENOPTERA.

(20) In Latin, *Chrysidida*.(22) In Latin, *Apida*.(24) In Latin, *Ichneumonida*.

V. EQUAL-WINGED INSECTS¹, with the mouth in form of a sucker, bent under the breast, comprehending,

- | | |
|--|---------------------------------|
| 1. Scale Insects ² . | 4. Chermes ⁵ . |
| 2. Plant Lice ³ , many without wings. | 5. Frog-hoppers ⁶ . |
| 3. Pierce Flies ⁴ . | 6. Lantern Flies ⁷ . |
| | 7. Tree-hoppers ⁸ . |

WING-CASED INSECTS⁹,

DIVIDED INTO

I. CROSS-WINGED INSECTS¹⁰, with the first pair of wings leathery at the base, overlapping the second pair crossways, and having a sucker bent under the breast, comprehending,

- | | |
|------------------------------------|--|
| 1. Boat Flies ¹¹ . | 6. Tree Bugs ¹⁶ . |
| 2. Water Scorpions ¹² . | 7. Common Bugs ¹⁷ : the Bed Bug, without wings. |
| 3. Water Measurers ¹³ . | |
| 4. Mask Bugs ¹⁴ . | |
| 5. Bank Bugs ¹⁵ . | |

II. LEATHER-WINGED INSECTS¹⁸, with the first pair of wings leathery throughout, overlapping at the edges only, and covering the second pair, which are folded lengthwise; comprehending,

- | | |
|----------------------------------|-----------------------------|
| 1. Cockroaches ¹⁹ . | 3. Locusts ²¹ . |
| 2. Mole Crickets ²⁰ . | 4. Crickets ²² . |

(1) In Latin, HOMOPTERA.

(2) In Latin, *Aphidæ*.

(3) In Latin, *Psyllidæ*.

(4) In Latin, *Fulgoridæ*.

(5) In Latin, ELYTROPTERA.

(6) In Latin, *Notonectidæ*.

(7) In Latin, *Hydrometridæ*.

(8) In Latin, *Coreidæ*.

(9) In Latin, *Cimicidæ*.

(10) In Latin, *Blattidæ*.

(11) In Latin, *Locustidæ*.

(12) In Latin, *Coccidæ*.

(13) In Latin, *Thripidæ*.

(14) In Latin, *Cercopicidæ*.

(15) In Latin, *Cicadidæ*.

(16) In Latin, HEMIPTERA.

(17) In Latin, *Nepidæ*.

(18) In Latin, *Reduidæ*.

(19) In Latin, *Pentatomidæ*.

(20) In Latin, ORTHOPTERA.

(21) In Latin, *Achetidæ*.

(22) In Latin, *Gryllidæ*.

III. HORN-CASED INSECTS¹, with the wing-cases, or first pair of wings, more or less horny, in some short, and some long, having the second pair of wings folded up, when at rest, beneath them; comprehending,

- | | |
|---|--|
| 1. Earwigs ² . In this I follow M. Straus-Dürckheim. | 16. Lady Birds ¹⁷ . |
| 2. Omalians ³ . | 17. Cassidians ¹⁸ . |
| 3. Stenians ⁴ . | 18. Chrysomelians ¹⁹ . |
| 4. Staphylinians ⁵ . | 19. Galerucians ²⁰ . |
| 5. Tachiporians ⁶ . | 20. Lily Beetles ²¹ . |
| 6. Pselaphians ⁷ . | 21. Leptureans ²² . |
| 7. Scydmanians ⁸ . | 22. Longhorned Beetles ²³ . |
| 8. Notoxians ⁹ . | 23. Salpingians ²⁴ . |
| 9. Blister Beetles ¹⁰ . | 24. Weevils ²⁵ . |
| 10. Cardinal Beetles ¹¹ . | 25. Bostrichians ²⁶ . |
| 11. Mordellians ¹² . | 26. Ptinians ²⁷ . |
| 12. Melandrians ¹³ . | 27. Tillians ²⁸ . |
| 13. Blapsians ¹⁴ . | 28. Telephorians ²⁹ . |
| 14. Tenebrions ¹⁵ . | 29. Glowworms ³⁰ . |
| 15. Endomychians ¹⁶ . | 30. Cebrians ³¹ . |
| | 31. Click Beetles ³² . |
| | 32. Buprestians ³³ . |

(1) In Latin, *COLEOPTERA*.

(2) In Latin, *Omalida*.

(3) In Latin, *Staphylinida*.

(4) In Latin, *Pselaphida*.

(5) In Latin, *Notoxida*.

(6) In Latin, *Cantharida*.

(7) In Latin, *Mordellida*.

(8) In Latin, *Blapsida*.

(9) In Latin, *Endomychida*.

(10) In Latin, *Coccinellida*.

(11) In Latin, *Cassidida*.

(12) In Latin, *Galerucida*.

(13) In Latin, *Lepturida*.

(14) In Latin, *Salpingida*.

(15) In Latin, *Bostrichida*.

(16) In Latin, *Ptinida*.

(17) In Latin, *Tillida*.

(18) In Latin, *Telephorida*.

(19) In Latin, *Cebrioida*.

(20) In Latin, *Elaterida*.

(21) In Latin, *Forficulida*.

(22) In Latin, *Stenida*.

(23) In Latin, *Tachyporida*.

(24) In Latin, *Scydmanida*.

(25) In Latin, *Cantharida*.

(26) In Latin, *Mordellida*.

(27) In Latin, *Blapsida*.

(28) In Latin, *Endomychida*.

(29) In Latin, *Cassidida*.

(30) In Latin, *Galerucida*.

(31) In Latin, *Lepturida*.

(32) In Latin, *Salpingida*.

(33) In Latin, *Bostrichida*.

(34) In Latin, *Ptinida*.

(35) In Latin, *Tillida*.

(36) In Latin, *Telephorida*.

(37) In Latin, *Cebrioida*.

(38) In Latin, *Elaterida*.

- | | |
|--|-----------------------------------|
| 33. Rose Chafers ¹ . | 46. Helophorians ¹⁴ . |
| 34. Cock Chafers ² . | 47. Limnians ¹⁵ . |
| 35. Dorr Beetles ³ . | 48. Parnians ¹⁶ . |
| 36. Stag Beetles ⁴ . | 49. Heterocerians ¹⁷ . |
| 37. Pill Beetles ⁵ . | 50. Whirlwigs ¹⁸ . |
| 38. Death-watch Beetles ⁶ . | 51. Water Beetles ¹⁹ . |
| 39. Engians ⁷ . | 52. Elaphrians ²⁰ . |
| 40. Nitulidians ⁸ . | 53. Bembidians ²¹ . |
| 41. Silphians ⁹ . | 54. Harpalians ²² . |
| 42. Scaphidians ¹⁰ . | 55. Carabians ²³ . |
| 43. Anisotamians ¹¹ . | 56. Scaritians ²⁴ . |
| 44. Sphæridians ¹² . | 57. Brachinians ²⁵ . |
| 45. Hydrophilians ¹³ . | 58. Cicindelians ²⁶ . |

TWO-WINGED INSECTS²⁷,

DIVIDED INTO

- I. PACT-WINGED FLIES²⁸, with only two wings, or none, and longish jaws; comprehending,
1. Bat Flies⁹. 2. Forest Flies³⁰.
- II. FAN-WINGED FLIES³¹, with the wings spreading somewhat like a fan; comprehending,
1. Bee Parasites³².

-
- | | |
|--------------------------------------|--------------------------------------|
| (1) In Latin, <i>Cetonidæ</i> . | (2) In Latin, <i>Melolonthidæ</i> . |
| (3) In Latin, <i>Scarabæidæ</i> . | (4) In Latin, <i>Lucanidæ</i> . |
| (5) In Latin, <i>Byrrhidæ</i> . | (6) In Latin, <i>Dermestidæ</i> . |
| (7) In Latin, <i>Engidæ</i> . | (8) In Latin, <i>Nitulidæ</i> . |
| (9) In Latin, <i>Silphidæ</i> . | (10) In Latin, <i>Scaphididæ</i> . |
| (11) In Latin, <i>Anisatomidæ</i> . | (12) In Latin, <i>Sphærididæ</i> . |
| (13) In Latin, <i>Hydrophilidæ</i> . | (14) In Latin, <i>Helophoridæ</i> . |
| (15) In Latin, <i>Limneidæ</i> . | (16) In Latin, <i>Parnidæ</i> . |
| (17) In Latin, <i>Heteroceridæ</i> . | (18) In Latin, <i>Gyrinidæ</i> . |
| (19) In Latin, <i>Dyticidæ</i> . | (20) In Latin, <i>Elaphridæ</i> . |
| (21) In Latin, <i>Bembididæ</i> . | (22) In Latin, <i>Harpalidæ</i> . |
| (23) In Latin, <i>Carabidæ</i> . | (24) In Latin, <i>Scaritidæ</i> . |
| (25) In Latin, <i>Brachinidæ</i> . | (26) In Latin, <i>Cicindelidæ</i> . |
| (27) In Latin, <i>Anelytra</i> . | (28) In Latin, HOMOLOPTEA. |
| (29) In Latin, <i>Nycteribidæ</i> . | (30) In Latin, <i>Hippoboscidæ</i> . |
| (31) In Latin, RHIPTEA. | (32) In Latin, <i>Stylopidæ</i> . |

III. POISERS¹, with poisers instead of the second pair of wings, and the mouth in form of a sucker; comprehending,

- | | |
|-----------------------------------|----------------------------------|
| 1. Common Flies ² . | 10. Bee Flies ¹¹ . |
| 2. Gad Flies ³ . | 11. Clegs ¹² . |
| 3. Conopians ⁴ . | 12. Mydasians ¹³ . |
| 4. Stomoxians ⁵ . | 13. Rhagions ¹⁴ . |
| 5. Wasp Flies ⁶ . | 14. Dolichoceans ¹⁵ . |
| 6. Wood Flies ⁷ . | 15. Empians ¹⁶ . |
| 7. Chameleon Flies ⁸ . | 16. Asilians ¹⁷ . |
| 8. Acrocerians ⁹ . | 17. Crane Flies ¹⁸ . |
| 9. Anthracians ¹⁰ . | 18. Gnats ¹⁹ . |

II. WINGLESS INSECTS²⁰,

[N.B.—Bed bugs, timber lice, worker ants, white ants, and some female moths, do not rank in this division, though they have no wings. We cannot make nature bend to our systems.]

DIVIDED INTO

I. LEAPING INSECTS²¹, with the hind legs formed for leaping; and comprehending,

1. Fleas²².

II. SPRINGING INSECTS²³, with the tail bristles formed for springing; and comprehending,

1. Spring Tails²⁴.

- | | |
|-------------------------------------|------------------------------------|
| (1) In Latin, DIPTERA. | (2) In Latin, <i>Muscidae</i> . |
| (3) In Latin, <i>Estridae</i> . | (4) In Latin, <i>Conopidae</i> . |
| (5) In Latin, <i>Stomoxyda</i> . | (6) In Latin, <i>Syrphidae</i> . |
| (7) In Latin, <i>Xylophagida</i> . | (8) In Latin, <i>Stratiomyda</i> . |
| (9) In Latin, <i>Acrocerida</i> . | (10) In Latin, <i>Anthracida</i> . |
| (11) In Latin, <i>Bombylida</i> . | (12) In Latin, <i>Tabanida</i> . |
| (13) In Latin, <i>Mydasida</i> . | (14) In Latin, <i>Rhagionida</i> . |
| (15) In Latin, <i>Dolichopida</i> . | (16) In Latin, <i>Empida</i> . |
| (17) In Latin, <i>Asilida</i> . | (18) In Latin, <i>Tipulida</i> . |
| (19) In Latin, <i>Culicida</i> . | (20) In Latin, APTERA. |
| (21) In Latin, THYSAURA. | (22) In Latin, <i>Pulicida</i> . |
| (23) In Latin, ANOPLURA. | (24) In Latin, <i>Podurida</i> . |

III. BLOODSUCKERS¹, with no members formed for leaping; comprehending,

1. Bird Lice².

2. Common Lice³.

It may be remarked that I have chiefly attended, in the preceding outline, to British insects.

(1) In Latin, *PARASITA*.

(2) In Latin, *Nirmidæ*.

(3) In Latin, *Pediculidæ*.

INDEX.

A

Acetabulum, 38
Achetidæ, 119
Acrida viridissima, 40, 45
Acroceridæ, 122
Aculeus, 61
Alæ, 37
Alæ anteriores, 47
Alæ posteriores, 55
Alulæ, 55
 ANELYTRA, 117, 121
Anisatomidæ, 121
 ANOPLURA, 122
Antefurca, 13
Antennæ, 10, 25
Antennulæ, 31
Anthracidæ, 122
Anus, 68
Apex anterior, 52
Apex inferior, 52
Aphidæ, 119
Apidæ, 118
Apodemæ articulationis, 20
Apodemæ insertionis, 20
 APTERA, 117, 122
Arcus, 22
Arcus tergi, 22
Arcus ventris, 22
Area, 51

Area exterior, 52
Area inferior, 52
Area media, 52
Area superior, 52
 ARTICULATA, 116
Asilidæ, 122
Aurelia, 108
Auriculo-ventriculariæ, 74

B

Basis, 52
Bembididæ, 121
Blapsidæ, 120
Blattidæ, 120
Bombycina, 118
Bombylidæ, 122
Bostrichidæ, 120
Brachinidæ, 121
Bulbus, 25
Buprestidæ, 120
Byrrhidæ, 121

C

Cæcum, 68
Calcaria, 38
Cantharidæ, 120
Caput, 9
Carabidæ, 121

Cardia, 66
Cardo, 30
Cassidæ, 120
Cauda, 53, 59
Cebrionidæ, 120
Cellulæ basilares, 52
Cellulæ exteriores, 52
Cellulæ mediæ, 52
Cerambycidæ, 120
Cercopicidæ, 119
Cetonidæ, 121
Chrysalis, 108
Chrysididæ, 118
Chrysomelidæ, 120
Cicadidæ, 119
Cicindelidæ, 121
Ciliæ, 53
Cimicidæ, 119
Clypeus, 29
Cocææ, 15
Coccidæ, 119
Coccinellidæ, 121
 COLEOPTERA, 120
Collum, 9
Conopidæ, 122
Coreidæ, 119
Corium, 5
Costa, 50
Costa anterior, 50
Costa inferior, 51
Costa interna, 51
Costa media, 50
Costa mediastina, 51
Costulæ marginales, 51
Coxa, 38
Cranium, 10
Crioceridæ, 120
Cubitus, 50
Culicidæ, 122
Curculionidæ, 120
Cuticula, 5
Cutis vera, 5
Cynipidæ, 118

D

Dermestidæ, 121
 DIPTERA, 122
Dolichopidæ, 122
Duodenum, 68
Dyticidæ, 121

E

Echinus, 65
Elaphridæ, 121
Elateridæ, 120
Elytra, 53
 ELYTROPTERA, 117, 119
Empidæ, 122
Endomychidæ, 120
Engidæ, 121
Entomology, 1
Entothorax, 13
Ephemeridæ, 118
Epidemæ, 48
Epidemæ articulationis, 20
Epidemæ insertionis, 20
Epimera, 15
Episterna, 15
Epistoma, 10
Eruca, 103
Excrementa, 68

F

Fæces, 68
Faliscus, 66
Fauces, 63
Fascia, 53
Femur, 38
Forceps, 59
Forficulidæ, 120
Formicidæ, 118
Frænnum, 18
Frænnum alæ, 56
Fulcrum, 34
Fulgoridæ, 119
Funiculus, 23
Furca, 13

G

- Galea*, 31
Galerucidæ, 120
Geometrina, 118
Gryllidæ, 119
Gyrinidæ, 121

H

- Halteres*, 56
Harpalidæ, 121
Helophorida, 121
HEMIPTERA, 119
Hesperidæ, 117
Heteroceridæ, 121
Hippoboscidæ, 121
HOMOLOPTERA, 121
HOMOPTERA, 119
Hydrophilidæ, 121
HYMENOPTERA, 118
Hydrometridæ, 119

I

- Ichneumonidæ*, 118
Imago, 109
Incunabulum, 107
Ingluvies, 65
INSECTA, 1
Intestina, 67
Intestinum tenue, 68
INVERTERRATA, 4
Irroratio, 53
Ischium primum, 16
Ischium secundum, 16

L

- Labium*, 29, 31
LAMBENTIA, 33
Lampyridæ, 120
Larva, 106
LEPIDOPTERA, 117
Lepisma saccharinum, 29

- Leptoceridæ*, 118
Lepturidæ, 120
Libellulidæ, 118
Limneidæ, 121
Lingula, 32
Lobus, 30
Locustidæ, 119
Lucanidæ, 121
Lycanidæ, 117

M

- Macula*, 53
Mælandridæ, 120
Mandibulæ, 29
MANDUCANTIA, 29
Margo anterior, 52
Margo exterior, 52
Margo inferior, 52
Maxillæ, 30
Melolonthidæ, 121
Mentum, 31
Mesonotum, 17
Mesosternum, 13
Mesostethium, 13
Mesothorax, 12
Metanotum, 17
Metasternum, 13
Metathorax, 12
Mordellidæ, 120
Muscidæ, 122
Mydasidæ, 122
Myrmeleonidæ, 118

N

- Nepidæ*, 119
Neuræ, 49
Neuroptera, 118
Nirmidæ, 123
Nitulidæ, 121
Noctuina, 118
Notonectidæ, 119
Notoxidæ, 120

Notum, 16
Nycteribidæ, 121
Nymphalidæ, 117

O.

Ocellus, 53
Œsophagus, 65
Œstridæ, 122
Omalidæ, 120
Omium, 18
 ORTHOPTERA, 119
Os, 29
Os hyoideum, 34
Ovarium, 97
Ovipositores, 61

P.

Palpi, 31
Palpi externi, 31
Palpi interni, 31
Palpus labialis, 31
Panorpidæ, 118
Papiliones, 117
Papilionidæ, 117
Paraglossa, 32
Parapleura, 16
Parapter, 15
 PARASITA, 123
Parnidæ, 121
Pars dorsalis, 30
Pars intermaxillaris, 30
Patagium, 15
Pediculidæ, 123
Pedes, 36
Pentatomidæ, 119
Peristethium, 13
Peritrema, 16
Perlidæ, 118
Phalænæ, 118
Phalanges, 38
Pharynx, 63
Phryganidæ, 118

Pleura, 16
Plumulæ, 56
Poduredæ, 122
Politum, 6
Postfurca, 13
Prehensores, 103
Proboscis, 33
Præscutum, 18
Profurca, 13
Pronotum, 17
Propedes, 103
Prophragma, 18
Prothorax, 12
Prosternum, 13
Pselaphidæ, 120
Psocidæ, 118
Psyllidæ, 119
 PTEROTA, 117
Ptinidæ, 120
Pubis, 18
Pulicidæ, 122
Pupa, 108
Pupum, 108
Pylorus, 67
Pyrochroidæ, 120

R.

Radius, 50
Ramuli, 78
Ramus transversalis, 30
Rectum, 68
Reduvidæ, 119
Rete mucosum, 5
Reticulæ, 52
Rhagienidæ, 123
 RHIPIDPTERA, 123
Rostrum, 38
Rotula, 150

S.

Salpingidæ, 12
Scaphididæ, 121
Scapulæ, 15

Scarabæidæ, 121
Scaritidæ, 121
Scutellum, 18
Scutum, 18
Scydmenidæ, 120
Segmentum, 22
Sericteria, 63
Sialisteria, 63
Sirphidæ, 122
Silphidæ, 121
Speisesack, 65
Sphærididæ, 121
Sphingina, 118
Spicula, 61
Spiracula, 75
Spiracula simplicia, 76
Squama palpifera, 30
Staphylinidæ, 120
Statio, 42
Stenidæ, 120
Sternum, 13
Sternum anterius, 13
Stigma, 51
Stigma primum, 13
Stigmata, 75
Stipes, 30
Stomoxydæ, 122
Stratiomydæ, 122
Striga, 53
Stylopidæ, 121
Suctoria, 35

T

Tabanidæ, 122
Tachyporidæ, 120
Tarsus, 38
Tela colorata, 5
Telephoridæ, 120
Tenebrionidæ, 121

Tenthredinidæ, 118
Tergum prothoracis, 17
Tergum mesothoracis, 17
Tergum metathoracis, 17
Tergum thoracis, 16
 TETRAPTERA, 117
Thorax, 12
Thripidæ, 119
 THYSANURA, 122
Tibia, 38
Tillidæ, 120
Tinodidæ, 118
Tipula oleracea, 42
Tipulidæ, 122
Tortricina, 118
Torus, 25
Tracheæ, 50, 72, 78
Tracheæ tubulariæ, 78
Tracheæ vesiculariæ, 79
 TRICHOPTERA, 118
Trochanter, 38
Trochantina, 38
Trochlea, 23

V U

Valvulainterventricularia, 74
Unguis, 39
Ventriculus, 65
 VERTEBRATA, 4
Vesicula dorsalis, 70
Vesiculi biliarii, 67
Vespidæ, 118
Villi, 66

X

Xylophagidæ, 122

Z

Zygæna statices, 35





